

Integration by Parts

Using integration by parts one transforms an integral of a product of two functions into a simpler integral. Divide the initial function into two parts called u and dv (keep dx in dv part). Then apply the following rule.



Integration by parts:

$$\int u dv = uv - \int v du$$

You were successful in choosing u and dv initially if the resulting integral $\int v du$ is **simpler** than the initial integral. If it is not, go back and rethink your choice of u and dv .

Below are some hints on how to decompose the initial integral into $u dv$:

- Integrals with the product of a polynomial and e^{ax} . Try $u = \text{polynomial}$.
- Integrals with polynomial and $\sin ax$ (or $\cos ax$). Try $u = \text{polynomial}$.
- $\int e^{ax} \sin bx \, dx$ or $\int e^{ax} \cos bx \, dx$. You can start with $u = e^{ax}$. You will need to do integration by parts twice and will end up with your initial integral after the second time. Solve for your initial integral then.
- Integrals with logarithmic functions. Try $u = \text{logarithmic function}$.
- Integrals with inverse trigonometric functions. Try $u = \text{inverse trigonometric function}$.

The formula $\int u dv = uv - \int v du$ is really the product rule in disguise. To see why the integration by parts formula is true, start with the product rule $(uv)' = u'v + v'u$ that can be also written as $\frac{d}{dx}(uv) = \frac{du}{dx}v + \frac{dv}{dx}u$. Integrating the product rule with respect to x , we have that $\int \frac{d}{dx}(uv) dx = \int \frac{du}{dx}v dx + \int \frac{dv}{dx}u dx \Rightarrow uv = \int dv u + \int v du = \int v du + \int u dv$. Solve for the term $\int u dv$ on the right side and obtain that $uv - \int v du = \int u dv$.

For definite integrals, the formula becomes

$$\int_a^b u dv = uv \Big|_a^b - \int_a^b v du.$$

Practice Problems. Evaluate the integrals:

1.
$$\int x e^x dx$$

2.
$$\int x e^{2x} dx$$

3.
$$\int x^2 e^x dx$$

Hypothesize on the number of integration by parts you would need to evaluate $\int x^n e^x dx$ or $\int x^n \sin x dx$.

4.
$$\int_0^1 x e^{-x} dx$$

5.
$$\int 2x \sin 3x dx$$

6.
$$\int (2x + 5) \sin(2x + 5) dx$$

7.
$$\int e^x \sin x dx$$

8.
$$\int \ln x dx$$

9.
$$\int \frac{\ln x}{x^2} dx$$

10.
$$\int \frac{\ln(2x - 1)}{(2x - 1)^2} dx$$

11.
$$\int \tan^{-1} x dx$$

12.
$$\int 3 \sin^{-1} 2x dx$$

Solutions:

1. Following the first hint, start by $u = x$ and $dv = e^x dx$. Then $du = dx$ and $v = \int e^x dx = e^x$. Then use the formula for integration by parts and obtain $\int x e^x dx = \int u dv = uv - \int v du = x e^x - \int e^x dx$. Note that this last integral is simpler than the initial integral indicating that you are on the right path. The last integral is equal to e^x so your final answer is $x e^x - e^x + c$.

2. Following the first hint, start by $u = x$ and $dv = e^{2x} dx$. Then $du = dx$ and $v = \int e^{2x} dx$. To get v , you can use the substitution $w = 2x \Rightarrow dw = 2dx \Rightarrow \frac{dw}{2} = dx$ and so $v = \int e^{2x} dx = \frac{1}{2} \int e^w dw = \frac{1}{2} e^w = \frac{1}{2} e^{2x}$.

Then $\int x e^{2x} dx = \int u dv = uv - \int v du = \frac{1}{2} x e^{2x} - \frac{1}{2} \int e^{2x} dx$. Note that this last integral is simpler than the initial integral indicating that you are on the right path. In fact, the last integral is the same as the one you evaluated when finding v so it is equal to $\frac{1}{2} e^{2x}$. Thus, the initial integral is equal to $\frac{1}{2} x e^{2x} - \frac{1}{2} \int e^{2x} dx = \frac{1}{2} x e^{2x} - \frac{1}{2} \cdot \frac{1}{2} e^{2x} + c = \frac{1}{2} x e^{2x} - \frac{1}{4} e^{2x} + c$

3. Start by $u = x^2$ and $dv = e^x dx$. Then $du = 2x dx$ and $v = \int e^x dx = e^x$. Then you have $\int x^2 e^x dx = \int u dv = uv - \int v du = x^2 e^x - \int e^x 2x dx = x^2 e^x + \int 2x e^x dx$. For this last integral, you need to use integration by parts again. Take $u = 2x$ and $dv = e^x dx$ (alternatively, factor 2 out and take $u = x$). Then $du = 2dx$ and $v = \int e^x dx = e^x$ and so $\int 2x e^x dx = 2x e^x - \int 2e^x dx = 2x e^x - 2e^x$. This gives you the final answer $\int x^2 e^x dx = x^2 e^x - (2x e^x - 2e^x) + c = x^2 e^x - 2x e^x + 2e^x + c$.

4. Start by $u = x$ and $dv = e^{-x} dx$. Then $du = dx$ and $v = \int e^{-x} dx$. To get v , you can use the substitution $w = -x$ and obtain $v = -e^{-x}$. Then $\int x e^{-x} dx = -x e^{-x} + \int e^{-x} dx = -x e^{-x} - e^{-x}$. Substituting the bounds you get $(-x e^{-x} - e^{-x})|_0^1 = -e^{-1} - e^{-1} + 1 = 1 - \frac{2}{e} \approx .264$.

5. Following the second hint, you can start by $u = 2x$ and $dv = \sin 3x dx$. Then $du = 2dx$ and $v = \int \sin 3x dx$. To find v , use the substitution $w = 3x \Rightarrow dw = 3dx \Rightarrow \frac{dw}{3} = dx$ and so $v = \int \sin 3x dx = \frac{1}{3} \int \sin w dw = \frac{-1}{3} \cos w = \frac{-1}{3} \cos 3x$. Then you have $\int 2x \sin 3x dx = \frac{-2}{3} x \cos 3x - \frac{-2}{3} \int \cos 3x dx$. This last integral is similar to the one used to obtain v and it is equal to $\frac{1}{3} \sin 3x$. Thus the initial integral is equal to $\frac{-2}{3} x \cos 3x + \frac{2}{3} \cdot \frac{1}{3} \sin 3x = \frac{-2}{3} x \cos(3x) + \frac{2}{9} \sin(3x) + c$.

6. **Method (1)** Use substitution $w = 2x + 5$ first to simplify the integral. In this case, $dw = 2dx \Rightarrow \frac{dw}{2} = dx$. Obtain $\int \frac{1}{2} w \sin w dw$. Using the integration by parts with $u = \frac{1}{2} w$ and $dv = \sin w dw$ we obtain $\frac{-1}{2} w \cos w + \frac{1}{2} \sin w + c = \frac{-1}{2} (2x + 5) \cos(2x + 5) + \frac{1}{2} \sin(2x + 5) + c$.

Method (2) Use the integration by parts with $u = 2x + 5$ and $dv = \sin(2x + 5)$. In this case, $du = 2dx$ and $v = \int \sin(2x + 5) dx$. To find v , you can use substitution $w = 2x + 5$. In this case, $dw = 2dx \Rightarrow \frac{dw}{2} = dx$. Obtain $v = \frac{1}{2} \int \sin w dw = \frac{-1}{2} \cos w = \frac{-1}{2} \cos(2x + 5)$. So, the integral becomes $\frac{-1}{2} (2x + 5) \cos(2x + 5) - \int \frac{-1}{2} \cos(2x + 5) 2dx$. This last integral simplifies to $\int \cos(2x + 5) dx$ (factor the negative out), you can evaluate it similarly as when finding v and obtain $\frac{1}{2} \sin(2x + 5)$. So, the final answer is $\frac{-1}{2} (2x + 5) \cos(2x + 5) + \frac{1}{2} \sin(2x + 5) + c$.

7. Note that the third hint applies to this problem. Let us denote the initial integral by I . You can start by $u = e^x$ and $dv = \sin x dx$. Then $du = e^x dx$ and $v = \int \sin x dx = -\cos x$ so that $I = \int e^x \sin x dx = -e^x \cos x + \int e^x \cos x dx$. Use the integration by parts again for this last integral. With $u = e^x$ and $dv = \cos x dx$, you obtain $du = e^x dx$ and $v = \sin x$ so that $\int e^x \cos x dx = e^x \sin x - \int e^x \sin x dx$. This last integral is our initial integral that we denoted by I . Thus

$$I = -e^x \cos x + \int e^x \cos x dx = -e^x \cos x + e^x \sin x - \int e^x \sin x dx = -e^x \cos x + e^x \sin x - I.$$

Note that this gives you the equation $I = -e^x \cos x + e^x \sin x - I$. Solving for I gives you

$$2I = -e^x \cos x + e^x \sin x \Rightarrow I = \frac{1}{2} (-e^x \cos x + e^x \sin x)$$

So, the final answer is $I = \frac{-1}{2}e^x \cos x + \frac{1}{2}e^x \sin x + c$.

8. Following the fourth hint, start by $u = \ln x$ and $dv = dx$. Then $du = \frac{1}{x}dx$ and $v = \int dx = x$. Thus $\int \ln x dx = x \ln x - \int \frac{1}{x}x dx = x \ln x - \int dx = x \ln x - x + c$.

9. Start by $u = \ln x$ and $dv = \frac{1}{x^2}dx = x^{-2}dx$ so that $du = \frac{1}{x}dx$ and $v = \int x^{-2}dx = -x^{-1} = \frac{-1}{x}$. Then $\int \frac{\ln x}{x^2} dx = \frac{-\ln x}{x} - \int \frac{-1}{x} \frac{1}{x} dx = \frac{-\ln x}{x} + \int x^{-2} dx = \frac{-\ln x}{x} - \frac{1}{x} + c$.

10. **Method (1)** Use substitution $w = 2x - 1$ first to simplify the integral. In this case, $dw = 2dx \Rightarrow \frac{dw}{2} = dx$. Obtain $\frac{1}{2} \int \frac{\ln w}{w^2} dw$. Then use integration by parts with $u = \ln w$ and $dv = \frac{dw}{w^2}$. Then $du = \frac{1}{w} dw$ and $v = \int w^{-2} dw = -w^{-1} = -\frac{1}{w}$. So, the integral becomes $\frac{-1}{2w} \ln w - \frac{1}{2} \int \frac{1}{w} \frac{-1}{w} dw = \frac{-\ln w}{2w} + \frac{1}{2} \int \frac{1}{w^2} dw = \frac{-\ln w}{2w} - \frac{1}{2} w^{-1} = \frac{-\ln w}{2w} - \frac{1}{2w} + c = \frac{-\ln(2x-1)}{2(2x-1)} - \frac{1}{2(2x-1)} + c$.

Method (2) Use integration by parts with $u = \ln(2x - 1)$ and $dv = \frac{dx}{(2x-1)^2}$. In this case, $du = \frac{1}{2x-1} 2dx$ and $v = \int (2x - 1)^{-2} dx$. You can use substitution $w = 2x - 1$ to find v . In this case, $dw = 2dx \Rightarrow \frac{dw}{2} = dx$. Obtain $v = \frac{1}{2} \int w^{-2} dw = \frac{-1}{2} w^{-1} = \frac{-1}{2w} = \frac{-1}{2(2x-1)}$. So, the integral becomes $\frac{-1}{2(2x-1)} \ln(2x - 1) - \int \frac{2}{2x-1} \frac{-1}{2(2x-1)} dx = \frac{-\ln(2x-1)}{2(2x-1)} + \int \frac{1}{(2x-1)^2} dx$. This last integral is the same one you evaluated when finding v , and so it is $\frac{-1}{2(2x-1)}$. Thus, the final answer is $\frac{-\ln(2x-1)}{2(2x-1)} - \frac{1}{2(2x-1)} + c$.

11. Following the fourth hint, start by $u = \tan^{-1} x$ and $dv = dx$. Thus $du = \frac{1}{1+x^2} dx$ and $v = \int dx = x$. So, the integral becomes $x \tan^{-1} x - \int \frac{x}{1+x^2} dx$. Use the substitution $w = 1 + x^2$ for this last integral and obtain that $\int \frac{x}{1+x^2} dx = \int \frac{x}{w} \frac{dw}{2x} = \frac{1}{2} \ln |w| = \frac{1}{2} \ln(1 + x^2)$. So, the initial integral is equal to $x \tan^{-1} x - \frac{1}{2} \ln(1 + x^2) + c$.

12. You can apply two methods to this problem as well: (1) use the substitution $w = 2x$ to simplify first and then use the integration by parts, or (2) use the integration by parts with $u = \sin^{-1}(2x)$ right away.

Using method (1), the integral reduces to $\int \frac{3}{2} \sin^{-1} w \frac{dw}{2}$. Then, using the integration by parts with $u = \sin^{-1} w$, you will obtain $\frac{3}{2} w \sin^{-1} w + \frac{3}{2} \sqrt{1 - w^2} + c$. Thus, the final answer is $3x \sin^{-1} 2x + \frac{3}{2} \sqrt{1 - 4x^2} + c$.

Using method (2), you have that $u = \sin^{-1}(2x)$ and $dv = 3dx$ so that $du = \frac{1}{\sqrt{1-(2x)^2}} 2dx = \frac{2}{\sqrt{1-4x^2}} dx$ and $v = \int 3dx = 3x$. Thus the integral is $3x \sin^{-1} 2x - \int 3x \frac{2}{\sqrt{1-4x^2}} dx$. Evaluate this integral using the substitution $w = 1 - 4x^2$ and obtain $\int 6x \frac{1}{\sqrt{w}} \frac{dw}{-8x} = \frac{-3}{4} \int w^{-1/2} dw = \frac{-3}{4} 2w^{1/2} = -\frac{3}{2} \sqrt{1 - 4x^2}$. So, the final answer is $3x \sin^{-1} 2x + \frac{3}{2} \sqrt{1 - 4x^2} + c$.