

## Review for Exam 1

1. **Surfaces.** Describe the following surfaces.

- (a)  $x^2 + y^2 = 4$       (b)  $x^2 + y^2 + z^2 = 9$       (c)  $z = 3$   
(d)  $z = 6 - 3x - 2y$       (e)  $z = \sqrt{16 - x^2 - y^2}$       (f)  $z = x^2 + y^2$   
(g)  $z = \sqrt{x^2 + y^2}$

2. **Review of Vectors.**

- (a) Let  $\vec{a} = \langle 3, 4, 0 \rangle$  and  $\vec{b} = \langle -1, 4, 2 \rangle$ . Find  $|\vec{a}|$ ,  $2\vec{a} + 3\vec{b}$ ,  $3\vec{a} - 2\vec{b}$ . Find the normalization of  $\vec{a}$ .  
(b) Find  $\vec{a} \cdot \vec{b}$  and  $\vec{a} \times \vec{b}$  for the two vectors from previous problem. Determine if the vectors are parallel, orthogonal or neither.  
(c) Let  $\vec{a} = \langle 1, 0, 1 \rangle$  and  $\vec{b} = \langle 1, -1, 0 \rangle$ . Find  $\vec{a} \cdot \vec{b}$ . Find the projection of  $\vec{b}$  onto  $\vec{a}$ .

3. **Lines and Planes.**

- (a) Find an equation of the line through the point  $(-2, 4, 10)$  and parallel to the vector  $\langle 3, 1, -8 \rangle$ .  
(b) Find an equation of the line through the point  $(1, 0, 6)$  and perpendicular to the plane  $x + 3y + z = 5$ .  
(c) Find an equation of the line through the points  $(3, 1, -1)$  and  $(3, 2, -6)$ .  
(d) Find an equation of the plane through the point  $(6, 3, 2)$  and perpendicular to the vector  $\langle -2, 1, 5 \rangle$ .  
(e) Find an equation of the plane through the point  $(4, -2, 3)$  and parallel to the plane  $3x - 7z = 12$ .  
(f) Find an equation of the plane through the points  $(0, 1, 1)$ ,  $(1, 0, 1)$  and  $(1, 1, 0)$ .  
(g) Find an equation of the line of the intersection of the planes  $x + y - z = 0$  and  $2x - 5y - z = 1$ .

4. **Curves in Space.**

- (a) Consider the curve  $x = \cos t$   $y = \sin t$   $z = t$ . Find an equation of the tangent line to the curve at the point where  $t = 0$ . Find the length of the curve from  $t = 0$  to  $t = 1$ .  
(b) Let  $C$  be the curve of intersection of the cylinder  $x^2 + y^2 = 1$  with the plane  $y + z = 2$ . Find parametric equations of this curve and an equation of the tangent line to the curve at the point where  $t = 0$  in parametrization. Using the calculator, estimate the length of the curve from  $t = 0$  to  $t = \pi/2$ .



## 8. Applications.

- (a) The pressure of 1 mole of an ideal gas is increasing at a rate of 0.05 kPa/s and the temperature is rising at a rate of 0.15 K/s. The pressure  $P$ , volume  $V$  and temperature  $T$  are related by the equation  $PV = 8.31T$ . Find the rate of change of the volume when the pressure is 20 kPa and temperature 320 K.
- (b) The number  $N$  of bacteria in a culture depends on temperature  $T$  and pressure  $P$  which depend on time  $t$  in minutes. Assume that 3 minutes after the experiment started, the pressure is increasing at a rate of 0.1 kPa/min and the temperature at a rate of 0.5 K/min. The number of bacteria changes at the rates of 3 bacteria per kPa and 5 bacteria per Kelvin. i) Find the rate at which the number of bacteria is increasing 3 minutes after the experiment started.  
ii) Assume that the rates of 3 bacteria per kPa and 5 bacteria per Kelvin are constant. If there is 300 bacteria initially when  $T = 305$  K and  $P = 102$  kPa, estimate the number of bacteria when  $T = 309$  K and  $P = 100$  kPa.
- (c) The number of flowers  $N$  in a closed environment depends on the amount of sunlight  $S$  that the flowers receive and the temperature  $T$  of the environment. Assume that the number of flowers changes at the rates  $N_S = 2$  and  $N_T = 4$ . i) If there are 100 flowers when  $S = 50$  and  $T = 70$ , estimate the number of flowers when  $S = 52$  and  $T = 73$ .  
ii) If the temperature depends on time as  $T(t) = 85 - \frac{8}{1+t^2}$  and the amount of sunlight decreases on time as  $S(t) = \frac{1}{t}$ , find the rate of change of the flower population at time  $t = 2$  days.
- (d) The temperature at a point  $(x, y)$  is  $T(x, y)$ , measured in degrees Celsius. A bug crawls so that its position after  $t$  seconds is given by  $x = \sqrt{1+t}$ ,  $y = 2 + \frac{1}{3}t$  where  $x$  and  $y$  are measured in centimeters. The temperature function satisfies  $T_x(2, 3) = 4$  and  $T_y(2, 3) = 3$ . Determine how fast the temperature rises on the bug's path after 3 seconds.
- (e) An object moves in the space away from its initial position so that after  $t$  hours it is at  $x = 2t$  and  $y = 2t^2 - 1$  and  $z = \sqrt{3t+1}$  miles from its initial position. Find the speed of that object 5 hours after it started moving. (Recall that the speed is the length of the tangent vector at a point.)

## Solutions

More detailed solutions of the problems can be found on the class handouts.

### 1. Surfaces.

- (a) Cylinder, base is a circle  $x^2 + y^2 = 4$  in  $xy$ -plane.      (b) Sphere, center at origin, radius 3.  
(c) Horizontal plane, passes  $(0, 0, 3)$ .      (d) Plane, passes  $(2, 0, 0)$ ,  $(0, 3, 0)$  and  $(0, 0, 6)$ .  
(e) Upper half-sphere, center is at origin, radius 4.      (f) Paraboloid, obtained by rotating a parabola  $z = y^2$  in  $zy$ -plane about  $z$ -axis.  
(g) Cone, obtained by rotating a line  $z = y$  in  $zy$ -plane about  $z$ -axis.

### 2. Review of Vectors.

- (a)  $|\vec{a}| = 5$ ,  $2\vec{a} + 3\vec{b} = \langle 3, 20, 6 \rangle$ ,  $3\vec{a} - 2\vec{b} = \langle 11, 4, -4 \rangle$ .  $\frac{\vec{a}}{|\vec{a}|} = \langle \frac{3}{5}, \frac{4}{5}, 0 \rangle$ .  
(b)  $\vec{a} \cdot \vec{b} = 13$ ,  $\vec{a} \times \vec{b} = \langle 8, -6, 16 \rangle$ .  $\vec{a}$  and  $\vec{b}$  are neither parallel nor orthogonal.  
(c)  $\vec{a} \cdot \vec{b} = 1$  The projection of  $\vec{b}$  onto  $\vec{a}$  is  $\langle \frac{1}{2}, 0, \frac{1}{2} \rangle$ .

### 3. Lines and Planes.

- (a)  $x = -2 + 3t$      $y = 4 + t$      $z = 10 - 8t$       (b)  $x = 1 + t$      $y = 3t$      $z = 6 + t$   
(c)  $x = 3$      $y = 1 + t$      $z = -1 - 5t$       (d)  $-2x + y + 5z = 1$   
(e)  $3x - 7z = -9$       (f)  $x + y + z = 2$       (g)  $x = 1 + 6t$      $y = t$      $z = 1 + 7t$

### 4. Curves in Space.

- (a) Tangent:  $x = 1, y = t, z = t$ . Length:  $\sqrt{2}$       (b) Tangent:  $x = 1, y = t, z = 2 - t$ .  
Length: 1.91  
(c) i)  $x = 3 \cos t, y = 3 \sin t, z = 1 - y^2 = 1 - 9 \sin^2 t$ . ii)  $(0, 3, -8)$  corresponds to  $t = \pi/2$ .  
Plugging  $\pi/2$  in derivative gives you  $\langle -3, 0, 0 \rangle$ . Tangent line:  $x = -3t$   $y = 3$   $z = -8$ . iii)  
 $(3, 0, 1)$  corresponds to  $t = 0$  and  $(0, 3, -8)$  to  $t = \pi/2$ . So, the bounds of integration are  
0 to  $\pi/2$ . The length is 10.48.  
(d) i)  $y = 4 \cos t, z = 4 \sin t, x = 8 - y^2 - z = 8 - 16 \cos^2 t - 4 \sin t$ . ii)  $(-8, -4, 0)$  corresponds  
to  $t = \pi$ . Plugging  $\pi$  in derivative gives you  $\langle 4, 0, -4 \rangle$ . Tangent line:  $x = 4t - 8$   $y = -4$   
 $z = -4t$ . iii)  $(4, 0, 4)$  corresponds to  $t = \pi/2$  and  $(-8, -4, 0)$  to  $t = \pi$ . So, the bounds of  
integration are  $\pi/2$  to  $\pi$ . The length is 14.515.  
(e) A set of parametric equations for the three curves in the intersection is  
 $x = 2 \cos t, y = 2 \sin t, z = 0$  with  $0 \leq t \leq \frac{\pi}{2}$ ,  
 $x = t, y = 0, z = 4 - t^2$  with  $0 \leq t \leq 2$ , and  
 $x = 0, y = t, z = 4 - t^2$  with  $0 \leq t \leq 2$ .

The three derivative vectors and length elements are

$$\begin{aligned}x' &= -2 \sin t, y' = 2 \cos t, z' = 0 \Rightarrow ds = \sqrt{4 \sin^2 t + 4 \cos^2 t} dt = \sqrt{4} dt = 2 dt \\x' &= 1, y' = 0, z' = -2t \Rightarrow ds = \sqrt{1 + 4t^2} dt, \text{ and} \\x' &= 0, y' = 1, z' = -2t \Rightarrow ds = \sqrt{1 + 4t^2} dt.\end{aligned}$$

The total length is  $\int_0^{\pi/2} 2 dt + \int_0^2 \sqrt{1 + 4t^2} dt + \int_0^2 \sqrt{1 + 4t^2} dt = \pi + 4.65 + 4.65 \approx 12.44$ .

5. Partial Derivatives.

- (a)  $z_x = 6x + 2y$ ,  $z_y = 2x - 10y$ ,  $z_{xx} = 6$ ,  $z_{xy} = z_{yx} = 2$ ,  $z_{yy} = -10$
- (b)  $z_x = e^x \sin y$ ,  $z_y = e^x \cos y$ ,  $z_{xx} = e^x \sin y$ ,  $z_{xy} = z_{yx} = e^x \cos y$ ,  $z_{yy} = -e^x \sin y$
- (c)  $z_x = 2axe^{x^2-xy} + ax^2e^{x^2-xy}(2x - y) = a(2x + 2x^3 - x^2y)e^{x^2-xy}$ ,  $z_y = ax^2e^{x^2-xy}(-x) = -ax^3e^{x^2-xy}$ . Then  $z_{xx} = a(2 + 6x^2 - 2xy)e^{x^2-xy} + a(2x + 2x^3 - x^2y)e^{x^2-xy}(2x - y)$  and  $z_{yy} = -ax^3e^{x^2-xy}(-x) = ax^4e^{x^2-xy}$ . Differentiating  $z_x$  with respect to  $y$  get  $z_{xy} = a(-x^2)e^{x^2-xy} + a(2x + 2x^3 - x^2y)e^{x^2-xy}(-x) = a(-x^2 - 2x^2 - 2x^4 + x^3y)e^{x^2-xy} = a(-3x^2 - 2x^4 + x^3y)e^{x^2-xy}$ . Alternatively, differentiating  $z_y$  with respect to  $x$  get  $z_{xy} = -3ax^2e^{x^2-xy} - ax^3e^{x^2-xy}(2x - y) = a(-3x^2 - 2x^4 + x^3y)e^{x^2-xy}$
- (d)  $z_x = \ln(xy^2) + 1$ ,  $z_y = 2x/y$ ,  $z_{xx} = 1/x$ ,  $z_{xy} = z_{yx} = 2/y$ ,  $z_{yy} = -2x/y^2$
- (e)  $z_x = -(y^2 + 2xz)/(2yz + x^2)$ ,  $z_y = -(2xy + z^2)/(2yz + x^2)$ . At  $(1, 1, 1)$ ,  $z_x = -1$ ,  $z_y = -1$ .
- (f)  $z_x = -(1 + \sin(x + y + z))/(-y + \sin(x + y + z))$  and  $z_y = -(-z + \sin(x + y + z))/(-y + \sin(x + y + z))$ . At  $(0, 1, -1)$ ,  $z_x = 1$  and  $z_y = 1$ .
- (g)  $z'(t) = (6x + 2y)(2t) + (2x - 10y)(-3t^2)$ ;  $z'(0) = 0$
- (h)  $z'(t) = (\ln(x + 2y) + x/(x + 2y))(-\sin t) + (2x/(x + 2y))(\cos t)$ ;  $z'(0) = 2$ .
- (i)  $z_s = (e^x y + y^2)t + (e^x + 2xy)2s$ ;  $z_t = (e^x y + y^2)s + (e^x + 2xy)2t$ ;  $z_s(1, 1) = 4e + 12$  and  $z_t(1, 1) = 4e + 12$
- (j)  $z_s = (2x + y)(-e^t \sin s) + x(e^t \cos s)$ ,  $z_t = (2x + y)(e^t \cos s) + x(e^t \sin s)$ ,  $z_s(\pi, 0) = 1$  and  $z_t(\pi, 0) = 2$

6. Tangent plane.

- (a)  $8x + 10y - z = 9$  (b)  $e^3y - z = e^3$
- (c)  $F_x = 2x, F_y = 4y, F_z = 6z$ . At  $(4, -1, 1)$  this produces vector  $\langle 8, -4, 6 \rangle$ . The tangent plane is  $4x - 2y + 3z = 21$ .
- (d)  $F_x = y^2 + 2xz, F_y = 2xy + z^2, F_z = 2yz + x^2$ . At  $(1, 1, 1)$  this produces vector  $\langle 3, 3, 3 \rangle$ . The tangent plane  $x + y + z = 3$ .
- (e)  $F_x = 1 + \sin(x + y + z), F_y = -z + \sin(x + y + z), F_z = -y + \sin(x + y + z)$ . At  $(0, 1, -1)$  this produces vector  $\langle 1, 1, -1 \rangle$ . The tangent plane is  $x + y - z = 2$ .

7. Linear Approximation. (a)  $f(2.02, 3.1) \approx 5.38$  (b)  $f(.9, 1.99) \approx 2.92$  (c)  $f(1.95, 1.08) \approx 2.847$  (d)  $f(6.9, 2.06) \approx -0.28$

8. Applications.

- (a) -.27 liter per second
- (b) i) Since  $P'(3) = 0.1$ ,  $T'(3) = 0.5$ ,  $N_P = 3$ , and  $N_T = 5$ , and  $N'(t) = N_T T' + N_P P'$  We have that  $N'(3) = 5 \cdot 0.5 + 3 \cdot 0.1 = 2.8$  bacteria/minute. ii) Using linear approximation formula,  $N(T, P) \approx N(T_0, P_0) + N_T \cdot (T - T_0) + N_P \cdot (P - P_0) \Rightarrow N(309, 100) \approx N(305, 102) + 5(309 - 305) + 3(100 - 102) = 300 + 5(4) + 3(-2) = 314$  bacteria.
- (c) i)  $100 + 2(52 - 50) + 4(73 - 70) = 116$  flowers. ii)  $N' = N_S S' + N_T T' = 2 \frac{-1}{4} + 4 \frac{32}{25} = 4.62$  flowers/day.
- (d) 2 degrees Celsius per second (e) Speed = 20.1 miles per hour.