

Derivatives and Integrals of Trigonometric and Inverse Trigonometric Functions

Trigonometric Functions.

Recall that

$$\begin{aligned} \text{if } y = \sin x, \text{ then } y' &= \cos x \quad \text{and} \\ \text{if } y = \cos x, \text{ then } y' &= -\sin x. \end{aligned}$$

Thus,

$$\begin{aligned} \int \sin x \, dx &= -\cos x + c \quad \text{and} \\ \int \cos x \, dx &= \sin x + c. \end{aligned}$$

The derivatives and integrals of the remaining trigonometric functions can be obtained by expressing these functions in terms of sine or cosine using the following identities: $\tan x = \frac{\sin x}{\cos x}$, $\cot x = \frac{\cos x}{\sin x}$, $\sec x = \frac{1}{\cos x}$, $\csc x = \frac{1}{\sin x}$.

Example 1. Find the derivative of $\tan x$. Simplify your answer.

Solution. Using the formula $\tan x = \frac{\sin x}{\cos x}$ and the quotient rule, obtain $\frac{d \tan x}{dx} = \frac{\cos x \cos x - (-\sin x) \sin x}{\cos^2 x} = \frac{\cos^2 x + \sin^2 x}{\cos^2 x} = \frac{1}{\cos^2 x}$ or $\sec^2 x$.

Inverse Trigonometric Functions. The function $\sin x$ passes horizontal line test for $-\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$ so it has an inverse. The inverse function is denoted by $\sin^{-1} x$ or $\arcsin x$. Since the range of $\sin x$ on $[-\frac{\pi}{2}, \frac{\pi}{2}]$ is $[-1, 1]$, the interval $[-1, 1]$ is the domain of $\sin^{-1} x$. We also have the following cancellation rule.

$$\sin(\sin^{-1} x) = x \quad \text{for } -1 \leq x \leq 1 \quad \text{and} \quad \sin^{-1}(\sin x) = x \quad \text{for } -\frac{\pi}{2} \leq x \leq \frac{\pi}{2}$$

Similarly, one obtains $\cos^{-1} x$ on $[0, \pi]$ is the inverse of $\cos x$ for $0 \leq x \leq \pi$ and the analogous cancellation rule holds. The function $\tan^{-1} x$ on $(-\infty, \infty)$ has the inverse $\tan x$ for $-\frac{\pi}{2} < x < \frac{\pi}{2}$. The inverses of other trigonometric functions can be obtained similarly.

Careful: when using notation $\sin^{-1} x$ do not mix this function up with $(\sin x)^{-1} = \frac{1}{\sin x}$.


When solving an equation of the form $\sin x = a$ for x where a is a number in $[-1, 1]$, the second cancellation formula implies that $x = \sin^{-1} a$ is one solution of this equation and that this value is in the interval $[-\frac{\pi}{2}, \frac{\pi}{2}]$. In many cases, we need more than this one solution, in particular, the second solution in interval $[0, 2\pi]$.

Using the trigonometric circle (or the graph of sine function) we can see that if

$$x_1 = \sin^{-1} a,$$

the second solution can be obtained as

$$x_2 = \pi - x_1 = \pi - \sin^{-1} a.$$



Similarly, when solving $\cos x = a$, in interval $[0, 2\pi]$ the first solution can be found as

$$x_1 = \cos^{-1} a$$

and the second solution can be obtained as


$$x_2 = -x_1 = -\cos^{-1} a.$$

For the equation $\tan x = a$, on $[0, 2\pi]$, the first solution can be found as

$$x_1 = \tan^{-1} a$$

and the second solution can be obtained as

$$x_2 = \pi + x_1 = \pi + \tan^{-1} a.$$



Derivatives of the Inverse Trigonometric Functions.

The formula for the derivative of $y = \sin^{-1} x$ can be obtained using the fact that the derivative of the inverse function $y = f^{-1}(x)$ is the reciprocal of the derivative $x = f(y)$.

$$y = \sin^{-1} x \Rightarrow x = \sin y \Rightarrow x' = \cos y \Rightarrow y' = \frac{1}{x'} = \frac{1}{\cos y} = \frac{1}{\cos(\sin^{-1} x)}.$$

To be able to simplify this last expression, one needs to represent $\cos y$ in terms of $\sin y$. This can be done using the trigonometric identity

$$\sin^2 y + \cos^2 y = 1 \Rightarrow \cos y = \sqrt{1 - \sin^2 y} \Rightarrow \cos(\sin^{-1} x) = \sqrt{1 - (\sin(\sin^{-1} x))^2} = \sqrt{1 - x^2}.$$

Thus, we obtain the formula for the derivative of $y = \sin^{-1} x$ to be

$$y' = \frac{1}{\sqrt{1 - x^2}}$$

Similarly, one obtains the following formulas.

$$\frac{d}{dx}(\tan^{-1} x) = \frac{1}{x^2 + 1} \qquad \frac{d}{dx}(\sec^{-1} x) = \frac{1}{x\sqrt{x^2 - 1}}$$

Differentiating the trigonometric identities $\sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$, $\tan^{-1} x + \cot^{-1} x = \frac{\pi}{2}$, and $\sec^{-1} x + \csc^{-1} x = \frac{\pi}{2}$, we obtain that the derivatives of $\cos^{-1} x$, $\cot^{-1} x$, and $\csc^{-1} x$ are negative of the derivatives of $\sin^{-1} x$, $\tan^{-1} x$, and $\sec^{-1} x$ respectively.

Example 2. Find the derivatives of the following functions.

(a) $y = \sin^{-1}(2x)$

(b) $y = x \tan^{-1} \sqrt{x}$

(c) $y = \sin^{-1} x^2 + \sqrt{1 - x^2}$

Solution. (a) Use the chain rule with $2x$ as the inner function and $\sin^{-1} x$ as the outer. The derivative of the outer with the inner function kept unchanged is $\frac{1}{\sqrt{1-(2x)^2}} = \frac{1}{\sqrt{1-4x^2}}$. The derivative of the inner function is 2 so the derivative of $y = \sin^{-1}(2x)$ is

$$y' = \frac{2}{\sqrt{1-4x^2}}.$$

(b) The function is a product of $f = x$ and $g = \tan^{-1} \sqrt{x}$. Use the chain rule for the derivative of g . Obtain that $f' = 1$ and $g' = \frac{1}{1+(\sqrt{x})^2} \cdot \frac{1}{2} x^{-1/2} = \frac{1}{2\sqrt{x}(1+x)}$. Hence the derivative of $y = x \tan^{-1} \sqrt{x}$ is

$$y' = f'g + g'f = 1 \cdot \tan^{-1} \sqrt{x} + \frac{1}{2\sqrt{x}(1+x)} \cdot x = \tan^{-1} \sqrt{x} + \frac{\sqrt{x}}{2(1+x)}.$$

(c) The function is a sum of two terms and you can differentiate term by term. Use the chain rule for the first term to get $\frac{1}{\sqrt{1-(x^2)^2}} \cdot 2x = \frac{2x}{\sqrt{1-x^4}}$. The derivative of the second term is $\frac{1}{2}(1-x^2)^{-1/2}(-2x) = -\frac{x}{\sqrt{1-x^2}}$. Hence the derivative of the function $y = \sin^{-1} x^2 + \sqrt{1-x^2}$ is

$$y' = \frac{2x}{\sqrt{1-x^4}} - \frac{x}{\sqrt{1-x^2}}.$$

Integrals producing inverse trigonometric functions. The above formulas for the the derivatives imply the following formulas for the integrals.

$$\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + c$$

$$\int \frac{1}{x^2+1} dx = \tan^{-1} x + c$$

$$\int \frac{1}{x\sqrt{x^2-1}} dx = \sec^{-1} x + c$$

Example 3. Evaluate the following integrals.

$$(a) \int \frac{1}{\sqrt{1-9x^2}} dx$$

$$(b) \int \frac{1}{9x^2+1} dx$$

Solution. (a) Note that the integrand matches the form $\frac{1}{\sqrt{1-u^2}}$ with $u^2 = 9x^2$. This produces the desired substitution $u^2 = 9x^2 \Rightarrow u = 3x$. Hence $du = 3dx \Rightarrow dx = \frac{du}{3}$. Thus we have that

$$\int \frac{1}{\sqrt{1-9x^2}} dx = \int \frac{1}{\sqrt{1-u^2}} \frac{du}{3} = \frac{1}{3} \int \frac{1}{\sqrt{1-u^2}} du = \frac{1}{3} \sin^{-1}(u) + c = \frac{1}{3} \sin^{-1}(3x) + c.$$

(b) Note that the integrand matches the form $\frac{1}{1+u^2}$ with $u^2 = 9x^2$. This produces the desired substitution $u^2 = 9x^2 \Rightarrow u = 3x$. Hence $du = 3dx \Rightarrow dx = \frac{du}{3}$. Thus we have that

$$\int \frac{1}{9x^2+1} dx = \int \frac{1}{u^2+1} dx \frac{du}{3} = \frac{1}{3} \int \frac{1}{1+u^2} du = \frac{1}{3} \tan^{-1}(u) + c = \frac{1}{3} \tan^{-1}(3x) + c.$$

Example 4. Compare the methods for evaluating the following integrals.

$$(a) \int \frac{x}{\sqrt{1-9x^2}} dx \qquad (b) \int \frac{1}{\sqrt{1-9x^2}} dx$$

Use your conclusion to determine the method for evaluating the integral

$$(c) \int \frac{x+1}{\sqrt{1-9x^2}} dx$$

Solution. (a) Although the denominator matches the one from the previous problem, the substitution $u = 3x$ would not work because of the x in the numerator. However, the substitution $u = 1 - 9x^2$ has $du = -18x dx \Rightarrow dx = \frac{du}{-18x}$ so the x term from du can cancel the x in the numerator after the substitution. So, you can evaluate this integral using the “standard” i.e. following the reasoning like on the handout “The Indefinite Integral – Review”. Thus we have that

$$\int \frac{x}{\sqrt{1-9x^2}} dx = \int \frac{x}{\sqrt{u}} \frac{du}{-18x} = \frac{-1}{18} \int u^{-1/2} du = \frac{-2}{18} u^{1/2} + c = \frac{-1}{9} \sqrt{1-9x^2} + c.$$

(b) This problem is the same as part (a) of Example 3.

(c) Separate the integral into a sum of two. The first matches part (a) and the second part (b).

Thus we have that

$$\int \frac{x+1}{\sqrt{1-9x^2}} dx = \int \frac{x}{\sqrt{1-9x^2}} dx + \int \frac{1}{\sqrt{1-9x^2}} dx = -\frac{1}{9} \sqrt{1-9x^2} + \frac{1}{3} \tan^{-1}(3x) + c.$$

Example 5. Evaluate the following integrals.

$$(a) \int \frac{1}{9+x^2} dx \qquad (b) \int \frac{1}{\sqrt{9-x^2}} dx$$

Solution. (a) To obtain the form $\frac{1}{1+u^2}$, factor 9 out of the denominator first. $\int \frac{1}{9+x^2} dx = \int \frac{1}{9(1+\frac{x^2}{9})} dx = \frac{1}{9} \int \frac{1}{1+\frac{x^2}{9}} dx$. Note that the integrand is of the form $\frac{1}{1+u^2}$, with $u^2 = \frac{x^2}{9}$. So, you can take $u = \frac{x}{3}$. Thus, $du = \frac{dx}{3} \Rightarrow 3du = dx$. Substitute and obtain $\frac{1}{9} \int \frac{1}{1+u^2} 3du = \frac{1}{3} \tan^{-1} u + c = \frac{1}{3} \tan^{-1}(\frac{x}{3}) + c$.

(b) Follow the same approach as in the previous problem: start by factoring 9 out of $9-x^2$. Obtain $\sqrt{9(1-\frac{x^2}{9})} = 3\sqrt{1-\frac{x^2}{9}}$. Thus the integral becomes $\int \frac{1}{3\sqrt{1-\frac{x^2}{9}}} dx = \frac{1}{3} \int \frac{1}{\sqrt{1-\frac{x^2}{9}}} dx$. Take $u^2 = \frac{x^2}{9}$ so that $u = \frac{x}{3}$. Thus, $du = \frac{dx}{3} \Rightarrow 3du = dx$. Substitute and obtain $\frac{1}{3} \int \frac{1}{\sqrt{1-u^2}} 3du = \sin^{-1} u + c = \sin^{-1}(\frac{x}{3}) + c$.

Example 6. Evaluate the integral

$$\int \frac{7}{5+3x^2} dx.$$

Solution. Follow the methods used in the previous problems: factor 7 out of the integral and then 5 out of the denominator.

$$\int \frac{7}{5+3x^2} dx = 7 \int \frac{1}{5(1+\frac{3x^2}{5})} dx = \frac{7}{5} \int \frac{1}{1+\frac{3x^2}{5}} dx.$$

The last integrand indicates that $u^2 = \frac{3x^2}{5}$ so that $u = \frac{\sqrt{3x}}{\sqrt{5}}$. Thus, $du = \frac{\sqrt{3}}{\sqrt{5}} dx \Rightarrow \frac{\sqrt{5}}{\sqrt{3}} du = dx$. Substitute and obtain $\frac{7}{5} \int \frac{1}{1+u^2} \frac{\sqrt{5}}{\sqrt{3}} du = \frac{7\sqrt{5}}{5\sqrt{3}} \int \frac{1}{1+u^2} du = \frac{7\sqrt{5}}{5\sqrt{3}} \tan^{-1} u + c = \frac{7\sqrt{5}}{5\sqrt{3}} \tan^{-1} \frac{\sqrt{3x}}{\sqrt{5}} + c$.

Practice Problems:

1. Find all solutions of the following equations on interval $[0, 2\pi]$.

- | | |
|------------------------------|-----------------------------------|
| (a) $\sin x = \frac{2}{5}$ | (b) $\cos^2 x = \frac{1}{4}$ |
| (c) $2 \tan x + 3 = 9$ | (d) $2 \cos^2 x + \cos x - 1 = 0$ |
| (e) $\sin x \cos x = \sin x$ | (f) $\sin x = \cos x$ |

2. Find the derivatives of the following functions.

- | | |
|--|---------------------------|
| (a) $y = \cot x^2$ | (b) $y = x^2 \cos x^2$ |
| (c) $y = \sin(3x + 2) \cos(2x - 3)$ | (d) $y = \sin^{-1}(2x)$ |
| (e) $y = \sin(ax) \cos(bx)$ where a and b are arbitrary constants. | |
| (f) $y = \sin^{-1}(x^4)$ | (g) $y = x^2 \cos^{-1} x$ |
| (h) $y = \tan^{-1}(e^x)$ | (i) $y = e^{\tan^{-1} x}$ |

3. Evaluate the following integrals.

- | | | |
|--|--|--|
| (a) $\int \cos(3x + 1) dx$ | (b) $\int x \sin x^2 dx$ | (c) $\int (9 + 2 \sin \frac{\pi x}{5}) dx$ |
| (d) $\int_0^1 \frac{1}{\sqrt{1-x^2}} dx$ | (e) $\int \frac{1}{\sqrt{1-4x^2}} dx$ | (f) $\int \frac{1}{4x^2+1} dx$ |
| (g) $\int \frac{1}{x^2+4} dx$ | (h) $\int \frac{x+1}{x^2+4} dx$ | (i) $\int \frac{3}{5x^2+8} dx$ |
| (j) $\int \frac{x+3}{5x^2+8} dx$ | (k) (Extra credit level) $\int \frac{x+6}{x^2+4x+13} dx$ | |

4. Find the area of the region between $y = \sin x$ and $y = \cos x$ for x in $[0, 2\pi]$.

Solutions.

- (a) $\sin x = \frac{2}{5} \Rightarrow x = \sin^{-1} \frac{2}{5}$ and $x = \pi - \sin^{-1} \frac{2}{5} \Rightarrow x \approx .411$ and $x \approx 2.73$ radians or 23.57 and 156.42 degrees.

(b) $\cos^2 x = \frac{1}{4} \Rightarrow \cos x = \pm \frac{1}{2} \Rightarrow x = \pm \cos^{-1} \frac{1}{2}, x = \pm \cos^{-1} \frac{-1}{2} \Rightarrow x = \pm \frac{\pi}{3}, x = \pm \frac{2\pi}{3}$. Converted to values in the interval $[0, 2\pi]$, the four solutions are $\frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}$ and $\frac{5\pi}{3}$ or 60, 120, 240, and 300 degrees.

(c) $2 \tan x + 3 = 9 \Rightarrow 2 \tan x = 6 \Rightarrow \tan x = 3 \Rightarrow x = \tan^{-1}(3)$ and $x = \pi + \tan^{-1}(3) \Rightarrow x \approx 1.25$ and $x \approx 4.39$ radians or 71.56 and 251.57 degrees.

(d) $2 \cos^2 x + \cos x - 1 = 0 \Rightarrow (2 \cos x - 1)(\cos x + 1) = 0 \Rightarrow \cos x = \frac{1}{2}, \cos x = -1 \Rightarrow x = \pm \cos^{-1} \frac{1}{2} = \pm \frac{\pi}{3}, x = \pm \cos^{-1}(-1) = \pm \pi$. Converted to values in the interval $[0, 2\pi]$, the solutions are $\frac{\pi}{3}, \pi,$ and $\frac{5\pi}{3}$ or 60, 180, and 300 degrees.

(e) $\sin x \cos x = \sin x \Rightarrow \sin x \cos x - \sin x = \sin x(\cos x - 1) = 0 \Rightarrow \sin x = 0, \cos x = 1$. The first equation has solutions 0 and π and the second just 0. Thus 0 and π are the solutions.

- (f) One way to solve this equation is to divide by $\cos x$ so that it reduces to $\tan x = 1$. Thus $x = \tan^{-1}(1) = \frac{\pi}{4}$ and $x = \pi + \tan^{-1}(1) = \frac{5\pi}{4}$. Another way is to convert \cos to sine (or vice versa). In this case the equation reduces to $\sin x = \sqrt{1 - \sin^2 x} \Rightarrow \sin^2 x = 1 - \sin^2 x \Rightarrow \sin^2 x = \frac{1}{2} \Rightarrow \sin x = \pm \frac{1}{\sqrt{2}}$. Be careful about the extraneous roots $\frac{3\pi}{4}$ and $\frac{7\pi}{4}$.
2. (a) Representing the function as $y = \cot x^2 = \frac{\cos x^2}{\sin x^2}$ and using the quotient rule with $f(x) = \cos x^2$ and $g(x) = \sin x^2$ and the chain for $f'(x) = -\sin x^2(2x)$ and $g'(x) = \cos x^2(2x)$ obtain that $y' = \frac{-\sin x^2(2x)\sin x^2 - \cos x^2(2x)\cos x^2}{\sin^2 x^2} = \frac{-2x(\sin^2 x^2 + \cos^2 x^2)}{\sin^2 x^2} = \frac{-2x}{\sin^2 x^2}$.
- (b) The product rule with $f(x) = x^2$ and $g(x) = \cos x^2$ and the chain for $g'(x) = -\sin x^2(2x)$ so that $y' = 2x \cos x^2 - \sin x^2(2x)(x^2) = 2x \cos x^2 - 2x^3 \sin x^2$.
- (c) Product and chain: $y' = 3 \cos(3x + 2) \cos(2x - 3) - 2 \sin(2x - 3) \sin(3x + 2)$.
- (d) Use the chain rule with the outer $\sin^{-1}(u)$ and the inner $2x$. The derivative of the outer with the inner unchanged is $\frac{1}{\sqrt{1-(2x)^2}}$ and the derivative of the inner is 2. Thus $y' = \frac{2}{\sqrt{1-4x^2}}$.
- (e) Product and chain: $y' = a \cos(ax) \cos(bx) - b \sin(bx) \sin(ax)$.
- (f) Use the chain rule with the outer $\sin^{-1}(u)$ and the inner x^4 . The derivative of the outer with the inner unchanged is $\frac{1}{\sqrt{1-(x^4)^2}}$ and the derivative of the inner is $4x^3$. Thus $y' = \frac{4x^3}{\sqrt{1-x^8}}$.
- (g) Use the product rule. $y' = 2x \cos^{-1} x - \frac{x^2}{\sqrt{1-x^2}}$.
- (h) Use the chain rule with the outer $\tan^{-1}(u)$ and the inner e^x . The derivative of the outer with the inner unchanged is $\frac{1}{1+(e^x)^2}$ and the derivative of the inner is e^x . Thus $y' = \frac{e^x}{1+e^{2x}}$.
- (i) Use the chain rule with the outer e^u and the inner $\tan^{-1} x$. The derivative of the outer with the inner unchanged is $e^u = e^{\tan^{-1} x}$ and the derivative of the inner is $\frac{1}{1+x^2}$. Thus $y' = \frac{e^{\tan^{-1} x}}{1+x^2}$.
3. (a) Use the substitution $u = 3x + 1$. The integral is $\frac{1}{3} \sin(3x + 1) + c$.
- (b) Use the substitution $u = x^2$. The integral is $-\frac{1}{2} \cos(x^2) + c$.
- (c) Use the substitution $u = \frac{\pi x}{5} \Rightarrow du = \frac{\pi}{5} dx \Rightarrow \frac{5du}{\pi} = dx$. The integral becomes $\int (9 + 2 \sin u) \frac{5du}{\pi} = \frac{5}{\pi} \int (9 + 2 \sin u) du = \frac{5}{\pi} (9u - 2 \cos u) + c = \frac{5}{\pi} (9 \frac{\pi x}{5} - 2 \cos \frac{\pi x}{5}) + c = 9x - \frac{10}{\pi} \cos \frac{\pi x}{5} + c$.
- (d) An antiderivative is $\sin^{-1} x$. Substituting the bounds, you obtain $\sin^{-1}(1) - \sin^{-1}(0) = \frac{\pi}{2}$.
- (e) The integrand has the form $\frac{1}{\sqrt{1-u^2}}$ that yields $\sin^{-1} u$ with $4x^2$ being u^2 . This tells you that $4x^2 = u^2$ (*careful*: not u but u^2). With $u^2 = 4x^2$, you can have $u = 2x$. Thus, $du = 2dx \Rightarrow \frac{du}{2} = dx$. Substitute and obtain $\int \frac{1}{\sqrt{1-u^2}} \frac{du}{2} = \frac{1}{2} \sin^{-1} u + c = \frac{1}{2} \sin^{-1}(2x) + c$.
- (f) The integrand has the form $\frac{1}{1+u^2}$ that yields $\tan^{-1} x$ with $4x^2$ being u^2 . This tells you that $4x^2 = u^2$. So, you can take $u = 2x$. Thus, $du = 2dx \Rightarrow \frac{du}{2} = dx$. Substitute and obtain $\int \frac{1}{u^2+1} \frac{du}{2} = \frac{1}{2} \tan^{-1} u + c = \frac{1}{2} \tan^{-1}(2x) + c$.
- (g) To obtain the form $\frac{1}{1+u^2}$, factor 4 out of the denominator first. $\int \frac{1}{x^2+4} dx = \int \frac{1}{4(\frac{x^2}{4}+1)} dx = \frac{1}{4} \int \frac{1}{\frac{x^2}{4}+1} dx$. Note that the integrand is of the form $\frac{1}{1+u^2}$, with $u^2 = \frac{x^2}{4}$. So, you can take $u = \frac{x}{2}$. Thus, $du = \frac{dx}{2} \Rightarrow 2du = dx$. Substitute and obtain $\frac{1}{4} \int \frac{1}{u^2+1} 2du = \frac{1}{2} \tan^{-1} u + c = \frac{1}{2} \tan^{-1}(\frac{x}{2}) + c$.

(h) Note that the function $\frac{x+1}{x^2+4}$ is the sum $\frac{x}{x^2+4} + \frac{1}{x^2+4}$. Integrate both terms. The first integral can be evaluated using the substitution $u = x^2 + 4 \Rightarrow du = 2xdx \Rightarrow \frac{du}{2x} = dx$. Thus, $\int \frac{x}{x^2+4} dx = \int \frac{x}{u} \frac{du}{2x} = \frac{1}{2} \ln|u| = \frac{1}{2} \ln(x^2 + 4)$. The second integral reduces to the previous problem with substitution $u = \frac{x}{2}$ and solution $\frac{1}{2} \tan^{-1}(\frac{x}{2})$. Thus, the final answer is $\frac{1}{2} \ln(x^2 + 4) + \frac{1}{2} \tan^{-1}(\frac{x}{2}) + c$.

(i) $\int \frac{3}{5x^2+8} dx = 3 \int \frac{1}{5x^2+8} dx = 3 \int \frac{1}{8(\frac{5}{8}x^2+1)} dx = \frac{3}{8} \int \frac{1}{\frac{5}{8}x^2+1} dx$. Take $u^2 = \frac{5x^2}{8}$ so that $u = \frac{\sqrt{5x}}{\sqrt{8}}$. Thus, $du = \frac{\sqrt{5}}{\sqrt{8}} dx \Rightarrow \frac{\sqrt{8}}{\sqrt{5}} du = dx$. Substitute and obtain $\frac{3}{8} \int \frac{1}{u^2+1} \frac{\sqrt{8}}{\sqrt{5}} du = \frac{3\sqrt{8}}{8\sqrt{5}} \int \frac{1}{u^2+1} du = \frac{3\sqrt{8}}{8\sqrt{5}} \tan^{-1} u + c = \frac{3\sqrt{8}}{8\sqrt{5}} \tan^{-1} \frac{\sqrt{5x}}{\sqrt{8}} + c$.

(j) The function $\frac{x+3}{5x^2+8}$ is the sum $\frac{x}{5x^2+8} + \frac{3}{5x^2+8}$. Integrate both terms. The first integral can be evaluated using the substitution $u = 5x^2 + 8 \Rightarrow du = 10xdx \Rightarrow \frac{du}{10x} = dx$. Thus, $\int \frac{x}{5x^2+8} dx = \int \frac{x}{u} \frac{du}{10x} = \frac{1}{10} \ln|u| = \frac{1}{10} \ln(5x^2 + 8)$.

The second integral is the same as the integral in part (i). We determined that it is equal to $\frac{3\sqrt{8}}{8\sqrt{5}} \tan^{-1} \frac{\sqrt{5x}}{\sqrt{8}}$. Thus, the final answer is $\frac{1}{10} \ln(5x^2 + 8) + \frac{3\sqrt{8}}{8\sqrt{5}} \tan^{-1} \frac{\sqrt{5x}}{\sqrt{8}} + c$.

(k) To apply the ideas for solving previous problems to this one, you want to complete the denominator to squares first (that is: to write the quadratic of the form $ax^2 + bx + c$ in the form $(px + q)^2 + r^2$). The denominator $x^2 + 4x + 13$ is equal to $x^2 + 2(2)x + 2^2 + 9$. Note that the first three terms are equal to $(x + 2)^2$. Thus, the denominator is equal to $(x + 2)^2 + 9$.

This tells you that you want to evaluate the integral $\int \frac{x+6}{x^2+4x+13} dx$ as the sum of two integrals. For the first you can take the whole denominator $(x+2)^2+9$ for u . This indicates how to decompose the numerator so that $du = 2(x+2)dx \Rightarrow \frac{du}{2(x+2)} = dx$ cancels the x -terms in the numerator. Thus $\int \frac{x+6}{x^2+4x+13} dx = \int \frac{x+2+4}{(x+2)^2+9} dx = \int \frac{x+2}{(x+2)^2+9} dx + \int \frac{4}{(x+2)^2+9} dx$.

For the first integral you obtain $\int \frac{1}{u} \frac{du}{2} = \frac{1}{2} \ln u = \frac{1}{2} \ln((x+2)^2 + 9)$.

Reduce the second integral to formula $\frac{1}{u^2+1}$ that yields $\tan^{-1}(u)$. Note that the denominator $(x+2)^2 + 9$ is equal to $9[(\frac{x+2}{3})^2 + 1]$. So, for the second integral, you can use the substitution $u = \frac{x+2}{3} \Rightarrow du = \frac{dx}{3} \Rightarrow 3du = dx$. This integral becomes $= \frac{4}{9} \int \frac{1}{(\frac{x+2}{3})^2+1} dx = \frac{4}{9} \cdot 3 \int \frac{1}{u^2+1} du = \frac{4}{3} \tan^{-1} u = \frac{4}{3} \tan^{-1} \frac{x+2}{3}$. Thus, the final answer is $\frac{1}{2} \ln((x+2)^2 + 9) + \frac{4}{3} \tan^{-1} \frac{x+2}{3} + c$.

4. Find intersections. The equation $\sin x = \cos x$ has solutions $x = \frac{\pi}{4}$ and $x = \frac{5\pi}{4}$ by problem 1 (f). Graph the functions and note that the area consists of 3 regions as in the graph on the right. $A = A_1 + A_2 + A_3 = \int_0^{\pi/4} (\cos x - \sin x) dx + \int_{\pi/4}^{5\pi/4} (\sin x - \cos x) dx + \int_{5\pi/4}^{2\pi} (\cos x - \sin x) dx = (\sin x + \cos x)|_0^{\pi/4} + (-\cos x - \sin x)|_{\pi/4}^{5\pi/4} + (\sin x + \cos x)|_{5\pi/4}^{2\pi} = 4\sqrt{2} \approx 5.657$.

