


Double Integrals over General Regions

Recall that the area of region D in xy -plane as in the figure below can be computed using the following integral.



$$A = \int_a^b (d(x) - c(x)) dx.$$

Now note that the integrand $d(x) - c(x)$ can be considered as the definite integral $\int_{c(x)}^{d(x)} dy$ since

$$\int_{c(x)}^{d(x)} dy = y \Big|_{c(x)}^{d(x)} = d(x) - c(x).$$

Thus, the area A can also be computed as the double integral

$$A = \int \int_D dx dy = \int_a^b \int_{c(x)}^{d(x)} dx dy.$$

General Double Integral. Using exactly the same argument to determine the bounds of integration, we can evaluate a double integral over region D as on the above figure of *any* function $z = f(x, y)$ of two variables defined on a region D . Arguing as above, the region D consist of all points (x, y) such that $a \leq x \leq b$ and $c(x) \leq y \leq d(x)$. Thus, the double integral can be evaluated as follows.

$$\int \int_D f(x, y) dx dy = \int_a^b \left(\int_{c(x)}^{d(x)} f(x, y) dy \right) dx.$$


In this case, the *order of integration matters* – the inner integral should be evaluated first. So, one needs to integrate the function $f(x, y)$ with respect to y first.

If $F(x, y)$ is the resulting antiderivative, the double integral reduces to a single integral as follows.

$$\int \int_D f(x, y) dx dy = \int_a^b \left(\int_{c(x)}^{d(x)} f(x, y) dy \right) dx =$$

$$\int_a^b (F(x, d(x)) - F(x, c(x))) dx.$$

The Volume. Suppose that f is positive on the region D . The **double integral** of f over D is the **volume** of the solid that lies above the region D and below the surface $z = f(x, y)$.




We have seen that the area of a region D in xy -plane can be computed as follows $A = \int \int_D dx dy$. This double integral can also be considered as the integral computing the volume under the horizontal plane $z = 1$ and above region D . This volume is equal to the product of the area of the base D and the height 1 so it is equal in size to the area of A .

Alternative Scenario. Assume that the region D is as on the figure on the right. In this case, bounds for y are constant and for x are not. In particular, $c \leq y \leq d$ and $a(y) \leq x \leq b(y)$. The area of the region D can be found as follows.

$$A = \int_c^d (b(y) - a(y)) dy = \int_c^d x \Big|_{a(y)}^{b(y)} dy =$$

$$\int_c^d \int_{a(y)}^{b(y)} dx dy \Rightarrow A = \int \int_D dx dy.$$




More generally, if $z = f(x, y)$ is a function of two variables defined over region D as above, we can evaluate a double integral over region D as follows.

$$\int \int_D f(x, y) dx dy = \int_c^d \left(\int_{a(y)}^{b(y)} f(x, y) dx \right) dy$$

Here the order of integration matters and the inner integral should be evaluated first (with respect to x). If $F(x, y)$ is the resulting antiderivative, the double integral reduces to a single integral as follows.

$$\int \int_D f(x, y) dx dy = \int_c^d \left(\int_{a(y)}^{b(y)} f(x, y) dx \right) dy =$$

$$\int_c^d (F(b(y), y) - F(a(y), y)) dy.$$



Practice problems.

1. Calculate the following double integrals.
 - a) $\int \int_D x^3 y^2 dx dy$ where D is given by $0 \leq x \leq 2, -x \leq y \leq x$.
 - b) $\int \int_D (x + 2y) dx dy$ where D is given by $0 \leq x \leq 1, 0 \leq y \leq x^2$.
 - c) $\int \int_D 2x dx dy$ where D is given by $0 \leq y \leq 1, y \leq x \leq e^y$.
2. Find the volume of the solid bounded by the plane $2x + 2y + z = 4$ in the first octant.
3. Find the volume of the solid under the surface $z = xy$ and above the triangle with vertices $(1, 1)$, $(4, 1)$ and $(1, 2)$.

- Evaluate the integral $\int \int_D y^3 dx dy$ where D is the triangular region with vertices $(0, 2)$, $(1, 1)$ and $(3, 2)$.
- Using a double integral, find the area of the region between parabola $y = x^2$ and the line $y = x$.


Solutions.

- $\int_0^2 \int_{-x}^x x^3 y^2 dx dy = \int_0^2 x^3 \frac{y^3}{3} \Big|_{-x}^x dx = \int_0^2 x^3 \frac{2x^3}{3} dx = \frac{2x^7}{21} \Big|_0^2 = \frac{256}{21}$
 - $\int_0^1 \int_0^{x^2} (x + 2y) dx dy = \int_0^1 (xy + y^2) \Big|_0^{x^2} dx = \int_0^1 (x^3 + x^4) dx = (\frac{x^4}{4} + \frac{x^5}{5}) \Big|_0^1 = \frac{9}{20}$
 - $\int_0^1 \int_y^{e^y} 2x dx dy$. Integrate with respect to x first. $\int_0^1 x^2 \Big|_y^{e^y} dy = \int_0^1 (e^{2y} - y^2) dy = 2.86$
- Solve the equation of the plane $2x + 2y + z = 4$ for z to obtain the function $z = 4 - 2x - 2y$ which you should integrate.

So, the volume is $\int \int_D (4 - 2x - 2y) dx dy$ where D is the region in xy -plane formed by the coordinate axes and the intersection of the given plane and the xy -plane. The xy -plane has the equation $z = 0$ and the plane $2x + 2y + z = 4$ intersects $z = 0$ at the line $2x + 2y + 0 = 4 \Rightarrow 2y = 4 - 2x \Rightarrow y = 2 - x$.


Alternatively, you can note that the given plane intersects the x -axis at $2x + 2(0) + 0 = 4 \Rightarrow x = 2$ and it intersects the y -axis at $2(0) + 2y + 0 = 4 \Rightarrow 2y = 4 \Rightarrow y = 2$. The line in xy -plane passing $(2,0)$, and $(0,2)$ has the slope $m = \frac{2-0}{0-2} = -1$ and the y -intercept $b = 2$ so it is $y = -x + 2$.

So, you are integrating over the shaded triangle on the figure on the right. The x -values are bounded by the values $x = 0$ and $x = 2$, and the y -values by the curves $y = 0$ and $y = 2 - x$.



So, the volume is given by $V = \int_0^2 \int_0^{2-x} (4 - 2x - 2y) dx dy = \int_0^2 (4y - 2xy - 2\frac{y^2}{2}) \Big|_0^{2-x} dx = \int_0^2 (4(2-x) - 2x(2-x) - (2-x)^2) dx = \int_0^2 (2-x)(4-2x-(2-x)) dx = \int_0^2 (2-x)(2-x) dx = \frac{-1}{3}(2-x)^3 \Big|_0^2 = \frac{8}{3} \approx 2.67$.

- The integral $\int \int_D xy dx dy$, where D is the region in xy -plane determined by the triangle with three given vertices, computes the volume. Graph the triangle first. Note that the x -values inside of the triangle are bounded by $x = 1$ and $x = 4$.




The upper y -bound is the line passing (1,2) and (4,1). This line has the slope $\frac{2-1}{1-4} = \frac{1}{-3}$. The point-slope equation gives you $y - 2 = \frac{-1}{3}(x - 1) \Rightarrow y = \frac{-1}{3}x + \frac{7}{3}$. The lower y -bound is determined by the line passing (1,1) and (4,1). This is the horizontal line $y = 1$.

So, the volume is $V = \int_1^4 \int_1^{\frac{-1}{3}x + \frac{7}{3}} xy dx dy = \int_1^4 x \frac{y^2}{2} \Big|_1^{\frac{-1}{3}x + \frac{7}{3}} dx = \int_1^4 x \left(\frac{(\frac{-1}{3}x + \frac{7}{3})^2}{2} - \frac{1}{2} \right) dx =$ (use calculator) $= \frac{31}{8}$.


4. Graph the triangle first. Note that the x -values inside of the triangle are bounded by $x = 0$ and $x = 3$. The upper y -bound is the horizontal line $y = 2$. The lower y -bound is the line passing (0, 2) and (1, 1) for $0 \leq x \leq 1$ and the line passing (1,1) and (3, 2) for $1 \leq x \leq 3$. So, you need to divide the region in two parts D_1 and D_2 where D_1 is the part of the triangle left of $x = 1$ and D_2 is the part of the triangle right from $x = 1$.

The line passing (0, 2) and (1, 1) has the slope $\frac{2-1}{0-1} = -1$ and y -intercept 2, so the equation is $y = -x + 2$. The line passing (1,1) and (3, 2) the slope $\frac{2-1}{3-1} = \frac{1}{2}$. The equation is $y - 1 = \frac{1}{2}(x - 1) \Rightarrow y = \frac{1}{2}x + \frac{1}{2}$.

The integral is $\int \int_{D_1} y^3 dx dy + \int \int_{D_2} y^3 dx dy = \int_0^1 \int_{-x+2}^2 y^3 dx dy + \int_1^3 \int_{\frac{1}{2}x + \frac{1}{2}}^2 y^3 dx dy = \int_0^1 \frac{y^4}{4} \Big|_{-x+2}^2 dx + \int_1^3 \frac{y^4}{4} \Big|_{\frac{1}{2}x + \frac{1}{2}}^2 dx = \int_0^1 (4 - \frac{(-x+2)^4}{4}) dx + \int_1^3 (4 - \frac{(\frac{1}{2}x + \frac{1}{2})^4}{4}) dx =$ (use calculator) $= \frac{147}{20}$.



Alternative (and shorter) way. Note that if you integrate with respect to y -first, you don't need to divide the region in two parts. The y -values inside of the triangle are bounded by $y = 1$ and $y = 2$. The lower x -bound is the line passing (0,2) and (1,1), given in terms of x . Thus $y = -x + 2 \Rightarrow x = -y + 2$. The upper x -bound is the line passing (1,1) and (3,2). $y = \frac{1}{2}x + \frac{1}{2} \Rightarrow 2y = x + 1 \Rightarrow x = 2y - 1$.



The integral is

$$\int_1^2 \int_{-y+2}^{2y-1} y^3 dx dy = \int_1^2 y^3 (2y - 1 - (-y + 2)) dy = \int_1^2 (3y^4 - 3y^3) dy = \frac{3y^5}{5} - \frac{3y^4}{4} \Big|_1^2 = \frac{147}{20}.$$

5. The area can be obtained as $\int \int_D dx dy$ where D is the region between the parabola and the line. The curves intersect at $x^2 = x \Rightarrow x(x - 1) = 0 \Rightarrow x = 0$ and $x = 1$. On this region the line $y = x$ is the upper curve and the parabola $y = x - x^2$ the lower. So the area is $A = \int_0^1 \int_{x-x^2}^x dx dy = \int_0^1 y \Big|_{x-x^2}^x dx = \int_0^1 (x - x^2) dx = (\frac{x^2}{2} - \frac{x^3}{3}) \Big|_0^1 = \frac{1}{6}$.