

## Review for the Final Exam

1. **Sequences.** Determine whether the following sequences are convergent or divergent. If they are convergent, find their limits.

(a)  $a_n = \left(\frac{1}{2}\right)^n$

(b)  $a_n = \frac{n+1}{2n-1}$

(c)  $a_n = \frac{n+3n^3}{4n^2+35n-7+2n^3}$ ,

(d)  $a_1 = 1, \quad a_{n+1} = \frac{1}{1+a_n}$

(e)  $a_0 = 0, \quad a_{n+1} = \sqrt{2 + a_n}$

- (f) When calculating the hydrogen ion concentration  $[\text{H}^+]$  in an acid-base system, the problem frequently boils down to finding the limit of a recursive sequence. For example, when hydrochloric acid HCl is dissolved in water, we have  $[\text{H}^+]_1 = C_{\text{HCl}}$  and

$$[\text{H}^+]_{n+1} = C_{\text{HCl}} + \frac{K_w}{[\text{H}^+]_n},$$

where  $C_{\text{HCl}}$  is the analytical concentration of HCl and  $K_w$  is the water's autoprotolysis constant that is equal to  $10^{-14}$  at 25 degrees Centigrade. If the analytical concentration of HCl  $C_{\text{HCl}}$  is equal to  $10^{-7}$ , find the hydrogen ion concentration  $[\text{H}^+]$  and its pH value.

2. **Sum of Series.** Find the sum of the following series.

(a)  $\sum_{n=2}^{\infty} \frac{2^{n+2}}{3^{n-1}}$

(b)  $\sum_{n=1}^{\infty} \frac{2^{2n}}{3^{n-1}}$

(c)  $2 - \frac{2}{3} + \frac{2}{9} - \frac{2}{27} + \frac{2}{81} - \dots$

(d)  $3 - \frac{3}{4} + \frac{3}{16} - \frac{3}{64} + \frac{3}{256} - \dots$

3. **Convergence of Series.** Determine whether the following series are convergent or divergent.

(a)  $\sum_{n=1}^{\infty} \frac{n^2}{13n^2+4n+5}$

(b)  $\sum_{n=0}^{\infty} \frac{3}{2^n}$

(c)  $\sum_{n=1}^{\infty} \frac{n^2}{n^2+5}$

(d)  $\sum_{n=1}^{\infty} \frac{1}{n^4}$

(e)  $1 + \frac{1}{8} + \frac{1}{27} + \frac{1}{64} + \frac{1}{125} + \dots$

(f)  $\sum_{n=1}^{\infty} \frac{1}{(n+3)^4}$

(g)  $1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots$

(h)  $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{2n}{4n+1}$

(i)  $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{2n}{4n^2+1}$

- (j)  $\frac{1}{2^1} + \frac{2}{2^2} + \frac{3}{2^3} + \frac{4}{2^4} + \dots$
- (k)  $\sum_{n=1}^{\infty} \frac{2^n}{n!}$
- (l)  $\sum_{n=1}^{\infty} \frac{2^n}{n^n}$

4. **Convergence of Power Series.** Find all the values of  $x$  for which the series converges.

- (a)  $\sum_{n=1}^{\infty} \frac{(x-1)^n}{n2^n}$
- (b)  $\sum_{n=1}^{\infty} \frac{x^n}{n!}$
- (c)  $\sum_{n=1}^{\infty} \frac{3^n x^n}{n+1}$
- (d)  $\sum_{n=1}^{\infty} \frac{(x-2)^{n+1}}{n3^n}$

5. **Power Series Expansion.** Find the power series expansion of the following functions centered at given point.

- (a)  $e^{2x}; \quad x = 0$
- (b)  $xe^{2x}; \quad x = 0$
- (c)  $\frac{1}{1-x^2}; \quad x = 0$
- (d)  $\frac{1}{1+x}; \quad x = 0$
- (e)  $\sin 3x; \quad x = 0$
- (f)  $\int_0^x e^{x^2} dx; \quad x = 0$

6. **Applications of Taylor Polynomials.**

- (a) Find the Taylor polynomial of the third degree centered at 0 for  $e^x$ .
- (b) Find the Taylor polynomial of the fourth degree centered at 0 for  $\sin x$ . Using the polynomial, evaluate  $\sin(0.2)$ .
- (c) Find the Taylor polynomial of the second degree centered at 0 for  $e^x \sin x$ . Using the polynomial, evaluate  $e^{1/2} \sin(1/2)$ .
- (d) If  $f(2) = 5$ ,  $f'(2) = 3$  and  $f''(2) = 1$ , approximate  $f(2.1)$ .
- (e) If  $f(2) = 5$ ,  $f'(2) = 3$ ,  $f''(2) = 1$ , and  $f'''(x) = 1/2$  approximate  $f(1.9)$ .
- (f) If  $f(1) = f'(1) = -1$ ,  $f''(1) = f'''(1) = 0$  and  $f^{iv}(1) = 2$ , approximate  $f(1.01)$ .
- (g) Approximate the function  $e^{\frac{hv}{kT}} - 1$  with the Taylor polynomial of the second degree in terms of  $v$ .
- (h) The magnitude of the electric field  $E$  of a single charge  $q$  can be described by  $E = \frac{kq}{r^2}$  where  $r$  is the distance between the field and the charge and  $k$  is a proportionality constant. If two opposite charges are at distance  $d$  from each other, the formula for the electric field changes to

$$E = \frac{kq}{(r-d)^2} - \frac{kq}{(r+d)^2} = \frac{kq}{r^2(1-\frac{d}{r})^2} - \frac{kq}{r^2(1+\frac{d}{r})^2}$$

Use the Taylor polynomial of the second degree of the function  $f(x) = \frac{1}{(1-x)^2}$  to show that the magnitude of the electric field  $E$  can be approximated as  $E \approx \frac{4kqd}{r^3}$ .

## 7. Lines and Planes.

- (a) Find an equation of the line through the point  $(1, 0, 6)$  and perpendicular to the plane  $x + 3y + z = 5$ .
- (b) Find an equation of the line through the points  $(3, 1, -1)$  and  $(3, 2, -6)$ .
- (c) Find an equation of the plane through the point  $(4, -2, 3)$  and parallel to the plane  $3x - 7z = 12$ .
- (d) Find an equation of the plane through the points  $(0, 1, 1)$ ,  $(1, 0, 1)$  and  $(1, 1, 0)$ .

## 8. Curves in Space.

- (a) Let  $C$  be the curve of intersection of the cylinder  $x^2 + y^2 = 1$  with the plane  $y + z = 2$ . Find an equation of the tangent line to the curve at the point where  $t = 0$ . Using the calculator, estimate the length of the curve from  $t = 0$  to  $t = \pi/2$ .
- (b) Consider the curve  $C$  which is the intersection of the surfaces

$$y^2 + z^2 = 16 \quad \text{and} \quad x = 8 - y^2 - z$$

- i) Find the parametric equations that represent the curve  $C$ .
- ii) Find the equation of the tangent line to the curve  $C$  at point  $(-8, -4, 0)$ .
- iii) Find the length of the curve from  $(4, 0, 4)$  to  $(-8, -4, 0)$ . Use the calculator to evaluate the integral that you are going to get.

## 9. Partial Derivatives. Find the indicated derivatives.

- (a)  $z = ax^2e^{x^2-xy}$  where  $a$  is a constant;  $z_x$ ,  $z_y$ ,  $z_{xx}$ ,  $z_{xy}$  and  $z_{yy}$ .
- (b)  $z = x \ln(xy^2)$ ;  $z_x$ ,  $z_y$ ,  $z_{xx}$ ,  $z_{xy}$  and  $z_{yy}$ .
- (c)  $xy^2 + yz^2 + zx^2 = 3$ ;  $z_x$  and  $z_y$  at  $(1, 1, 1)$ .
- (d)  $x - yz = \cos(x + y + z)$ ;  $z_x$  and  $z_y$  at  $(0, 1, -1)$ .

## 10. Tangent planes. Find the equation of the tangent plane to a given surface at a specified point.

- (a)  $z = y^2 - x^2$ , at  $(-4, 5, 9)$
- (b)  $x^2 + 2y^2 + 3z^2 = 21$ , at  $(4, -1, 1)$
- (c)  $xy^2 + yz^2 + zx^2 = 3$ ; at  $(1, 1, 1)$ .

## 11. Linear Approximation.

- (a) If  $f(1, 2) = 3$ ,  $f_x(1, 2) = 1$  and  $f_y(1, 2) = -2$ , approximate  $f(.9, 1.99)$ .
- (b) Find the linear approximation of  $z = \sqrt{20 - x^2 - 7y^2}$  at  $(2, 1)$  and use it to approximate the value at  $(1.95, 1.08)$ .

## 12. Applications.

- (a) The pressure of 1 mole of an ideal gas is increasing at a rate of 0.05 kPa/s and the temperature is rising at a rate of 0.15 K/s. The pressure  $P$ , volume  $V$  and temperature  $T$  are related by the equation  $PV = 8.31T$ . Find the rate of change of the volume when the pressure is 20 kPa and temperature 320 K.

- (b) The temperature at a point  $(x, y)$  is  $T(x, y)$ , measured in degrees Celsius. A bug crawls so that its position after  $t$  seconds is given by  $x = \sqrt{1+t}$ ,  $y = 2 + \frac{1}{3}t$  where  $x$  and  $y$  are measured in centimeters. The temperature function satisfies  $T_x(2, 3) = 4$  and  $T_y(2, 3) = 3$ . How fast is the temperature rising on the bug's path after 3 seconds?
- (c) The number of flowers  $N(S, T)$  in a closed environment depends on the amount of sunlight  $S$  that the flowers receive and the temperature  $T$  of the environment. Assume that  $N_S = 2$  and  $N_T = 4$ . i) Assume that there are 100 flowers when  $S = 50$  and  $T = 70$ . Use the linear approximation to estimate the number of flowers when  $S = 52$  and  $T = 73$ . ii) If the temperature depends on time as  $T(t) = 85 - 8/(1+t^2)$  and the amount of sunlight decreases on time as  $S = 1/t$  find the rate of change of the flower population  $N'(t)$  at time  $t = 2$  days.
- (d) Find the points on the surface  $z^2 = xy + 1$  that are closest to the origin.
- (e) Set up the equations for finding the dimensions of the rectangular box with the largest volume if the total surface area is  $64 \text{ cm}^2$ .
- (f) A cardboard box without a lid is to have volume of  $32,000 \text{ cm}^3$ . Set up the equations for finding the dimensions that minimize the amount of cardboard used.

13. **Maximum and Minimum Values.** Find the maximum and minimum values of  $f$ .

(a)  $z = 9 - 2x + 4y - x^2 - 4y^2$                       (b)  $z = x^2 + y^2 + x^2y + 4$

14. **Lagrange Multipliers.** Find the maximum and minimum values of  $f$  subject to the given constraint(s).

(a)  $f(x, y) = x^2 - y^2$ ;  $x^2 + y^2 = 1$                       (b)  $f(x, y, z) = 2x + 6y + 10z$ ;  $x^2 + y^2 + z^2 = 35$

15. **Double Integrals.**

- (a)  $\int \int_D (x + 2y) dx dy$  where  $D = \{ (x, y) \mid 0 \leq x \leq 1, 0 \leq y \leq x^2 \}$
- (b)  $\int \int_D 2x dx dy$  where  $D = \{ (x, y) \mid 0 \leq y \leq 1, y \leq x \leq e^y \}$
- (c)  $\int \int_D y^3 dx dy$  where  $D$  is the triangular region with vertices  $(0, 2)$ ,  $(1, 1)$  and  $(3, 2)$
- (d) Find the average value of the function  $f(x, y) = 4x$  on the region  $D$  between the parabolas  $y = x^2 - 2$  and  $y = 3x - x^2$ .

16. **The Volume.**

- (a) Find the volume of the solid bounded by the plane  $x + y + z = 1$  in the first octant.
- (b) Find the volume of the solid under the paraboloid  $z = x^2 + y^2$  and above the disk  $x^2 + y^2 \leq 9$ .
- (c) Find the volume of the solid above the cone  $z = \sqrt{x^2 + y^2}$  and below the paraboloid  $z = 2 - x^2 - y^2$ .
- (d) Find the volume of the solid enclosed by the paraboloids  $z = x^2 + y^2$  and  $z = 36 - 3x^2 - 3y^2$ .

17. **Surface Area. Parametric Surfaces**

- (a) Find the area of the part of the plane  $3x + 2y + z = 6$  that lies in the first octant.

- (b) Find the area of the part of the surface  $z = y^2 - x^2$  that lies between the cylinders  $x^2 + y^2 = 1$  and  $x^2 + y^2 = 4$ .
- (c) Find the equation of the tangent plane to the parametric surface  $x = u + v$ ,  $y = 3u^2$ ,  $z = u - v$  at the point  $(2, 3, 0)$ .
- (d) Find the surface area of the part of the cone  $z = \sqrt{x^2 + y^2}$  that lies between the cylinders  $x^2 + y^2 = 4$  and  $x^2 + y^2 = 9$ . Write down the parametric equations of the cone first. Then find the surface area using the parametric equations.
- (e) Using the parametric equations and formula for the surface area for parametric curves, show that the surface area of the cylinder  $x^2 + z^2 = 4$  for  $0 \leq y \leq 5$  is  $20\pi$ .

### 18. Triple Integrals and volume.

- (a)  $\int \int \int_E xy \, dx \, dy \, dz$  where  $E$  is the solid tetrahedron with vertices  $(0,0,0)$ ,  $(1, 0, 0)$ ,  $(0, 2, 0)$  and  $(0, 0, 3)$ .
- (b)  $\int \int \int_E 2 \, dx \, dy \, dz$  where  $E$  is the solid that lies between the cylinders  $x^2 + y^2 = 1$  and  $x^2 + y^2 = 4$  and between the  $xy$ -plane and the plane  $z = x + 2$ .
- (c)  $\int \int \int_E x^2 + y^2 + z^2 \, dx \, dy \, dz$  where  $E$  is the unit ball  $x^2 + y^2 + z^2 \leq 1$ .
- (d)  $\int \int \int_E z \, dx \, dy \, dz$  where  $E$  is the region between the spheres  $x^2 + y^2 + z^2 = 1$  and  $x^2 + y^2 + z^2 = 4$  in the first octant.
- (e) Find the volume of the ellipsoid  $\frac{x^2}{4} + \frac{y^2}{9} + \frac{z^2}{25} = 1$  by using the transformation  $x = 2u$ ,  $y = 3v$ ,  $z = 5w$ .

### 19. Line Integrals.

- (a)  $\int_C x y^4 \, ds$ ,  $C$  is the right half of the circle  $x^2 + y^2 = 16$ .
- (b)  $\int_C (xy + \ln x) \, dy$ ,  $C$  is the parabola  $y = x^2$  from  $(1,1)$  to  $(3,9)$ .
- (c)  $\int_C xy \, dx + (x - y) \, dy$ ,  $C$  consists of line segments from  $(0,0)$  to  $(2,0)$  and from  $(2,0)$  to  $(3,2)$ .
- (d)  $\int_C z^2 \, dx + y \, dy + 2y \, dz$ , where  $C$  consists of two parts  $C_1$  and  $C_2$ .  $C_1$  is the intersection of the cylinder  $x^2 + y^2 = 16$  and the plane  $z = 3$  from  $(0, 4, 3)$  to  $(-4, 0, 3)$ .  $C_2$  is a line segment from  $(-4, 0, 3)$  to  $(0, 1, 5)$ .
- (e) Find the work done by the force field  $\vec{F}(x, y, z) = (x + y^2, y + z^2, z + x^2)$  in moving an object along the curve  $C$  which is the intersection of the plane  $x + y + z = 1$  and the coordinate planes.
- (f) Find the work done by the force field  $\vec{F} = (-y^2, x, z^2)$  in moving an object along the curve  $C$  which is the intersection of the plane  $y + z = 2$  and the cylinder  $x^2 + y^2 = 1$ .

### 20. Potential. Independence of Path.

- (a) Check that  $\vec{f} = \langle x^3 y^4, x^4 y^3 + 2y \rangle$  is conservative, find its potential function and use it to evaluate  $\int_C \vec{f} \cdot d\vec{r}$  where  $C$  is  $x = \sqrt{t}$ ,  $y = 1 + t^3$ ,  $0 \leq t \leq 1$ .
- (b) Check that  $\vec{f} = \langle y, x + z, y \rangle$  is conservative, find its potential function and use it to evaluate  $\int_C \vec{f} \cdot d\vec{r}$  where  $C$  is any path from  $(2, 1, 4)$  to  $(8, 3, -1)$ .

- (c) Show that the line integral  $\int_C(2xy + z^2)dx + (x^2 + 2yz + 2)dy + (y^2 + 2xz + 3)dz$  where  $C$  is any path from  $(1, 0, 2)$  to  $(0, 1, 4)$ , is independent of path and evaluate it.

21. **Green's Theorem.** Evaluate the following integrals using Green's theorem.

- (a)  $\oint_C x^4 dx + xy dy$  where  $C$  is the triangle with vertices  $(0, 0)$ ,  $(0, 1)$ , and  $(1, 0)$ . Compute the integral also without using Green's Theorem.
- (b)  $\oint_C y^2 dx + 3xy dy$  where  $C$  is the boundary of the region between the circles  $x^2 + y^2 = 1$  and  $x^2 + y^2 = 4$  above  $x$ -axis.
- (c)  $\oint_C e^y dx + 2xe^y dy$  where  $C$  is the square with vertices  $(0, 0)$ ,  $(1, 0)$ ,  $(1, 1)$ , and  $(0, 1)$ .
- (d)  $\oint_C xy dx + 2x^2 dy$  where  $C$  is the line segment from  $(-2, 0)$  to  $(2, 0)$  and the upper half of the circle  $x^2 + y^2 = 4$ .

22. **Curl and Divergence.** Find curl and divergence of the following vector fields.

(a)  $\vec{f} = \langle xz, xyz, -y^2 \rangle$

(b)  $\vec{f} = \langle e^x \sin y, e^x \cos y, z \rangle$

## Solutions

More detailed solutions of the problems can be found on the class handouts.

- Sequences. (a) convergent, limit is 0 (b) convergent, limit is  $1/2$  (c) convergent, limit is  $3/2$  (d) convergent, find limit from the equation  $x = \frac{1}{1+x}$ , the limit is .618 (e) convergent, find limit from the equation  $x = \sqrt{2+x}$ , the limit is 2 (f) Find limit from the equation  $x^2 - 10^{-7}x - 10^{-14} = 0$ . The relevant solution is  $[H^+] = 1.618 \cdot 10^{-7}$  and pH= 6.7910.
- Sum of Series. (a)  $\frac{2^2}{3^{-1}} \sum_{n=2}^{\infty} \frac{2^n}{3^n} = \frac{12\frac{4}{9}}{\frac{1}{3}} = 16$ . (b)  $\sum_{n=1}^{\infty} 3 \left(\frac{4}{3}\right)^n$ .  $r = \frac{4}{3} > 1$  so the series is divergent. (c) sum= $3/2$  (d) sum =  $12/5$
- Convergence of Series.
  - Divergent by the Divergence Test
  - Geometric Series. Convergent because  $-1 < r = \frac{1}{2} < 1$ .
  - Divergent by the Divergence Test
  - $p$ -series. Convergent because  $4 > 1$
  - $p$ -series,  $p = 3$ . Convergent because  $3 > 1$
  - Convergent by the Integral Test
  - Note that the series is  $\sum_{n=1}^{\infty} (-1)^{n+1} \frac{1}{n}$ . Use the Alternating Series Test with  $b_n = \frac{1}{n}$ . The sequence  $b_n$  has limit 0 and is decreasing. Thus, the series is convergent.
  - Divergent by the Divergence Test

- (i) Convergent by the Alternating Series Test ( $b_n = \frac{2n}{4n^2+1}$  has limit 0 and is decreasing)
- (j) Note that the series is  $\sum_{n=1}^{\infty} \frac{n}{2^n}$ . Use the Ratio Test. The limit from the test is  $\frac{1}{2}$  which is less than 1 and so the series is convergent.
- (k) Convergent by the Ratio Test (limit from the test is 0 which is less than 1)
- (l) Convergent by the Root Test (limit from the test is 0 which is less than 1)
4. Convergence of Power Series. Series converges for (a)  $-1 \leq x < 3$  (b) All values of  $x$   
(c)  $-1/3 \leq x < 1/3$  (d)  $-1 \leq x < 5$ .
5. Power Series Expansion. (a)  $e^{2x} = \sum_{n=0}^{\infty} \frac{2^n x^n}{n!}$  (b)  $xe^{2x} = \sum_{n=0}^{\infty} \frac{2^n x^{n+1}}{n!}$  (c)  $\frac{1}{1-x^2} = \sum_{n=0}^{\infty} x^{2n}$  (d)  $\frac{1}{1+x} = \sum_{n=0}^{\infty} (-1)^n x^n$  (e)  $\sin 3x = 3x - \frac{3^3 x^3}{3!} + \frac{3^5 x^5}{5!} - \frac{3^7 x^7}{7!} + \dots = \sum_{n=0}^{\infty} \frac{(-1)^n 3^{2n+1} x^{2n+1}}{(2n+1)!}$ . (f)  $e^{x^2} = \sum_{n=0}^{\infty} \frac{x^{2n}}{n!} \Rightarrow \int_0^x e^{x^2} dx = \sum_{n=0}^{\infty} \frac{x^{2n+1}}{n!(2n+1)}$ .
6. Applications of Taylor Polynomials. (a)  $1+x+x^2/2+x^3/6$  (b)  $0+x+0-x^3/6+0 = x-x^3/6$ .  
 $\sin(0.2) \approx .1987$  (c)  $0+x+x^2 \cdot e^{1/2} \sin(1/2) \approx .75$  (d)  $f(2.1) \approx 5+3(.1)+1/2(.1)^2 = 5.305$   
(e)  $f(1.9) \approx 5+3(-.1)+(1/2)(-.1)^2+1/12(-.1)^3 = 4.705$  (f)  $f(1.01) \approx -1-1(.01)+2/24(.01)^4 = -1.00999 \approx -1.01$
- (g)  $f(v) = e^{\frac{hv}{kT}} - 1 \Rightarrow f'(v) = \frac{h}{kT} e^{\frac{hv}{kT}} \Rightarrow f''(v) = \frac{h^2}{k^2 T^2} e^{\frac{hv}{kT}}$ . Thus  $f(0) = 1 - 1 = 0$ ,  $f'(0) = \frac{h}{kT}$ ,  
 $f''(0) = \frac{h^2}{k^2 T^2}$ . So  $f(v) \approx \frac{hv}{kT} + \frac{h^2 v^2}{2k^2 T^2} = \frac{hv(2kT+hv)}{2k^2 T^2}$ .
- (h)  $\frac{1}{(1-x)^2} \approx -1-2x-3x^2$  and  $\frac{1}{(1+x)^2} \approx -1+2x-3x^2 \Rightarrow E = \frac{kq}{r^2} \left( -1 + 2\frac{d}{r} - 3\frac{d^2}{r^2} + 1 + 2\frac{d}{r} + 3\frac{d^2}{r^2} \right) = \frac{4kqd}{r^3}$ .
7. Lines and Planes. (a)  $x = 1+t$   $y = 3t$   $z = 6+t$  (b)  $x = 3$   $y = 1+t$   $z = -1-5t$   
(c)  $3x - 7z = -9$  (d)  $x + y + z = 2$
8. Curves in Space. (a) Tangent:  $x = 1$ ,  $y = t$ ,  $z = 2 - t$ . Length: 1.91 (b) i)  $y = 4 \cos t$ ,  
 $z = 4 \sin t$ ,  $x = 8 - y^2 - z = 8 - 16 \cos^2 t - 4 \sin t$ . ii)  $(-8, -4, 0)$  corresponds to  $t = \pi$ . Plugging  
 $\pi$  in derivative gives you  $\langle 4, 0, -4 \rangle$ . Tangent line:  $x = 4t - 8$   $y = -4$   $z = -4t$ . iii)  $(4, 0, 4)$   
corresponds to  $t = \pi/2$  and  $(-8, -4, 0)$  to  $t = \pi$ . The length is 14.515.
9. Partial Derivatives.
- (a)  $z_x = 2ax e^{x^2-xy} + ax^2 e^{x^2-xy}(2x-y) = a(2x+2x^3-x^2y)e^{x^2-xy}$ ,  $z_y = ax^2 e^{x^2-xy}(-x) = -ax^3 e^{x^2-xy}$ . Then  $z_{xx} = a(2+6x^2-2xy)e^{x^2-xy} + a(2x+2x^3-x^2y)e^{x^2-xy}(2x-y)$  and  $z_{yy} = -ax^3 e^{x^2-xy}(-x) = ax^4 e^{x^2-xy}$ . Differentiating  $z_x$  with respect to  $y$  get  $z_{xy} = a(-x^2)e^{x^2-xy} + a(2x+2x^3-x^2y)e^{x^2-xy}(-x) = a(-x^2-2x^2-2x^4+x^3y)e^{x^2-xy} = a(-3x^2-2x^4+x^3y)e^{x^2-xy}$ . Alternatively, differentiating  $z_y$  with respect to  $x$  get  $z_{xy} = -3ax^2 e^{x^2-xy} - ax^3 e^{x^2-xy}(2x-y) = a(-3x^2-2x^4+x^3y)e^{x^2-xy}$
- (b)  $z_x = \ln(xy^2) + 1$ ,  $z_y = 2x/y$ ,  $z_{xx} = 1/x$ ,  $z_{xy} = z_{yx} = 2/y$ ,  $z_{yy} = -2x/y^2$
- (c)  $z_x = -(y^2+2xz)/(2yz+x^2)$ ,  $z_y = -(2xy+z^2)/(2yz+x^2)$ . At  $(1, 1, 1)$ ,  $z_x = -1$ ,  $z_y = -1$ .
- (d)  $z_x = -(1+\sin(x+y+z))/(-y+\sin(x+y+z))$  and  $z_y = -(-z+\sin(x+y+z))/(-y+\sin(x+y+z))$ . At  $(0, 1, -1)$ ,  $z_x = 1$  and  $z_y = 1$ .
10. Tangent planes. (a)  $8x + 10y - z = 9$  (b)  $F_x = 2x, F_y = 4y, F_z = 6z$ . At  $(4, -1, 1)$  this produces vector  $\langle 8, -4, 6 \rangle$ . The tangent plane is  $4x - 2y + 3z = 21$ . (c)  $F_x = y^2 + 2xz, F_y = 2xy + z^2, F_z = 2yz + x^2$ . At  $(1, 1, 1)$  this produces vector  $\langle 3, 3, 3 \rangle$ . The tangent plane  $x+y+z = 3$ .

11. Linear Approximation. (a)  $f(.9, 1.99) \approx 2.92$  (b)  $f(1.95, 1.08) \approx 2.847$
12. Applications.
- (a) -.27 liter per second (b) 2 degrees Celsius per second
- (c) i)  $100 + 2(52 - 50) + 4(73 - 70) = 116$  flowers. ii)  $N_t = N_S S_t + N_T T_t = 2(-1/4) + 432/25 = 4.62$  flowers/day. (d)  $(0, 0, \pm 1)$
- (e) Equations:  $yz - 2\lambda y - 2\lambda z = 0$ ,  $xz - 2\lambda x - 2\lambda z = 0$ ,  $xy - 2\lambda x - 2\lambda y = 0$ ,  $2xy + 2yz + 2xz = 64$ . If solved, the equations would yield:  $x = y = z = 4\sqrt{6}/3$  cm.
- (f) Equations:  $y + 2z - yz\lambda = 0$ ,  $x + 2z - xz\lambda = 0$ ,  $2x + 2y - xy\lambda = 0$ ,  $xyz = 32,000$ . (If solved, the equations would yield: square base of side  $x = y = 40$  cm, height  $z = 20$  cm.)
13. Maximum and Minimum Values. (a) Maximum  $f(-1, 1/2) = 11$  (b) Minimum  $f(0, 0) = 4$ , saddle points  $(\pm\sqrt{2}, -1)$
14. Lagrange Multipliers. (a) Max.  $f(\pm 1, 0) = 1$ , min.  $f(0, \pm 1) = -1$  (b) Max.  $f(1, 3, 5) = 70$ , min.  $f(-1, -3, -5) = -70$
15. Double Integrals. (a)  $\frac{9}{20}$  (b) 2.86 (c)  $\frac{147}{20}$  (d) mass = 6, center of mass =  $(\frac{3}{4}, \frac{3}{2})$
16. The Volume. (a)  $\frac{1}{6}$  (b)  $\frac{81\pi}{2}$  (c)  $\frac{5\pi}{6}$  (d)  $162\pi$
17. Surface Area. Parametric Surfaces.
- (a)  $3\sqrt{14}$  (b) 30.85 (c) Plane  $3x - y + 3z = 3$
- (d) Parameterization:  $x = r \cos t$ ,  $y = r \sin t$ ,  $z = \sqrt{x^2 + y^2} = \sqrt{r^2} = r$ . The length of the cross product is  $\sqrt{2}r$ . The surface area is  $5\pi\sqrt{2}$ .
- (e) Parameterization:  $x = 2 \cos t$ ,  $y = y$ ,  $z = 2 \sin t$ . Bounds:  $0 \leq t \leq 2\pi$ ,  $0 \leq y \leq 5$ . Length of the cross product is 2. Thus the double integral is  $2\pi \cdot 5 \cdot 2 = 20\pi$ .
18. Triple Integrals and volume. (a)  $\frac{1}{10}$  (b)  $12\pi$  (c)  $\frac{4\pi}{5}$  (d)  $\frac{15\pi}{16}$  (e)  $40\pi$
19. Line Integrals. (a) 1638.4 (b) 102.68 (c)  $17/3$  (d)  $\int_{C_1} = -44$ ,  $\int_{C_2} = 67.83$ . So,  $\int_C = 67.83 - 44 = 23.83$
- (e) Let  $C_1$  be a line from  $(1, 0, 0)$  to  $(0, 1, 0)$ ,  $C_2$  a line from  $(0, 1, 0)$  to  $(0, 0, 1)$  and  $C_3$  a line from  $(0, 0, 1)$  to  $(1, 0, 0)$ . Find that  $\int_{C_1} = \frac{-1}{3}$ ,  $\int_{C_2} = \frac{-1}{3}$ , and  $\int_{C_3} = \frac{-1}{3}$ . Thus  $\int_C = \int_{C_1} + \int_{C_2} + \int_{C_3} = -1$ .
- (f)  $C$  has parametrization  $x = \cos t$ ,  $y = \sin t$ ,  $z = 2 - y = 2 - \sin t$ ,  $0 \leq t \leq 2\pi$ .  $\int_C = \int_C -y^2 dx + x dy + z^2 dz = \int_0^{2\pi} \sin^3 t dt + \cos^2 t dt + (2 - \sin t)^2 \cos t dt = \pi$ .
20. Potential. Independence of Path. (a)  $F = \frac{1}{4}x^4 y^4 + y^2 + c$ ,  $\int_C = F(1, 2) - F(0, 1) = 7$  (b)  $F = xy + yz + c$ ,  $\int_C = F(8, 3, -1) - F(2, 1, 4) = 15$  (c)  $F = x^2 y + z^2 x + y^2 z + 2y + 3z + c$ ,  $\int_C = F(0, 1, 4) - F(1, 0, 2) = 8$ .
21. Green's Theorem. (a)  $\frac{1}{6}$  (b)  $\frac{14}{3}$  (c)  $e - 1$  (d) 0
22. Curl and Divergence. (a)  $\text{div } \vec{f} = z + xz$ ,  $\text{curl } \vec{f} = \langle -y(x + 2), x, yz \rangle$  (b)  $\text{div } \vec{f} = 1$ ,  $\text{curl } \vec{f} = \langle 0, 0, 0 \rangle$