

Review for Exam 4

1. (a) Show that the relation \sim used to define \mathbb{Z} and given on $\mathbb{N} \times \mathbb{N}$ by

$$(k, l) \sim (m, n) \Leftrightarrow k + n = l + m$$

is an equivalence relation.

- (b) Show that the integer $[(0, 0)]$ is the identity for addition (i.e. show that $[(m, n)] + [(0, 0)] = [(m, n)]$ and $[(0, 0)] + [(m, n)] = [(m, n)]$).
- (c) For any integer $[(m, n)]$, show that its additive inverse is $[(n, m)]$ (i.e. show that $[(m, n)] + [(n, m)] = [(0, 0)]$ and $[(n, m)] + [(m, n)] = [(0, 0)]$).
- (d) Show that $[(n, m)] = [(0, m - n)]$ if $m \geq n$ and $[(n, m)] = [(n - m, 0)]$ if $m < n$ for any natural numbers m and n .
2. The relation \leq on \mathbb{Z} matches the familiar order of integers when $[(m, n)]$ is shortened to $m - n$. Rearrange the integer numbers below, if needed, so that the elements in the new list are non-decreasing.

$$[(6, 3)], [(1000, 1005)], [(6, 8)], [(57, 56)], [(56, 58)]$$

3. If $[(m, n)] \in \mathbb{Q}$ is such that $m \neq 0$, show that $[(m, n)] \cdot [(n, m)] = [(1, 1)]$ and $[(n, m)] \cdot [(m, n)] = [(1, 1)]$.
4. The relation \leq on \mathbb{Q} matches the familiar order of integers when $[(m, n)]$ is shortened to $\frac{m}{n}$. Rearrange the rational numbers below, if needed, so that the elements in the new list are non-decreasing.

$$[(5, 15)], [(50, -100)], [(15, 10)], [(20, -10)], [(-10, 20)]$$

5. Show that \mathbb{Q} has no zero divisors, that is $ab = 0 \Rightarrow a = 0$ or $b = 0$ for any $a, b \in \mathbb{Q}$. You can assume that \mathbb{Z} has no zero divisors (that is $mn = 0 \Rightarrow m = 0$ or $n = 0$ for $m, n \in \mathbb{Z}$).
6. Find the limit of the following recursive sequences.

(a) $a_{n+1} = \sqrt{2 + a_n}, \quad a_0 = 0$

(b) $a_{n+1} = \frac{1}{1+a_n}, \quad a_0 = 1.$

7. Show that the following pairs of sets are in a bijective correspondence. You can assume the existence of any of the bijective correspondences from the formula sheet.

(a) $(3, 5) \cup [8, 9)$ and $(7, \infty)$

(b) $(3, 5] \cup [0, 9) \cup [7, \infty)$ and $(-\infty, 1]$

(c) $\bigcup_{n \in \mathbb{N} - \{0\}} (-n, n)$ and $(0, 1)$

(d) $\bigcap_{n \in \mathbb{N}} [0, n + 1)$ and \mathbb{R}

(e) $\bigcup_{n \in \mathbb{N}} (-\infty, -n)$ and $(1, \infty)$

8. Represent the following decimal numbers as quotients of two integer numbers.

(a) 0.222222...

(b) 0.27272727...

(c) 1.2345454545...

9. Determine the moduli and the arguments given the following complex numbers in algebraic forms: $-3i$, $\sqrt{2} - \sqrt{2}i$, $-\sqrt{3} + i$, $-2 - i$.
10. Determine the real and imaginary parts of the complex numbers given by their moduli and arguments: $\theta = \frac{-\pi}{2}, r = 5$; $\theta = \frac{5\pi}{6}, r = 2$; $\theta = \frac{-2\pi}{3}, r = 3$.
11. Determine the n -th power of the given complex numbers and given n . Express your answers in algebraic form. (a) $z = -\sqrt{3} + i, n = 4$; (b) $z = -2 - i, n = 6$.
12. Find all solutions of the following equations.
- (a) $z^5 + 32 = 0$ (b) $z^5 - 32 = 0$
(c) $z^4 = 3 + 3i$ (d) $z^4 = 3 - 3i$
13. Using the definitions of the complex-valued trigonometric functions $\sin z = \frac{1}{2i}(e^{iz} - e^{-iz})$ and $\cos z = \frac{1}{2}(e^{iz} + e^{-iz})$, show the identities below.
- (a) $e^{iz} = \cos z + i \sin z$ (b) $\sin^2 z + \cos^2 z = 1$
(c) $\sin^2 z = \frac{1}{2}(1 - \cos(2z))$ (d) $\cos^2 z = \frac{1}{2}(1 + \cos(2z))$

Solutions

1. (a) *Reflexivity*. We need to show that $(k, l) \sim (k, l)$ for any nonnegative integers k and l .

$$\begin{aligned}(k, l) \sim (k, l) &\Leftrightarrow k + l = l + k \quad (\text{by the definition of } \sim) \\ &\Leftrightarrow k + l = k + l \quad (\text{by commutativity of } +)\end{aligned}$$

Since $k + l = k + l$ is true, we have that $(k, l) \sim (k, l)$ is also true.

Symmetry. Assume that $(k, l) \sim (m, n)$ for some $k, l, m, n \in \mathbb{N}$ and show that $(m, n) \sim (k, l)$.

$$\begin{aligned}(k, l) \sim (m, n) &\Leftrightarrow k + n = l + m \quad (\text{by the definition of } \sim) \\ &\Leftrightarrow m + l = n + k \quad (\text{by commutativity of } + \text{ and symmetry of } =) \\ &\Leftrightarrow (m, n) \sim (k, l) \quad (\text{by the definition of } \sim)\end{aligned}$$

Transitivity. Assume that $(k, l) \sim (m, n)$ and that $(m, n) \sim (o, p)$ and show that $(k, l) \sim (o, p)$.

$$\begin{aligned}(k, l) \sim (m, n) \wedge (m, n) \sim (o, p) &\Leftrightarrow k + n = l + m \wedge m + p = n + o \quad (\text{by the definition of } \sim) \\ &\Rightarrow k + n + m + p = l + m + n + o \quad (\text{by adding the equations}) \\ &\Leftrightarrow k + p = l + o \quad (\text{by cancelling } n + m) \\ &\Leftrightarrow (k, l) \sim (o, p) \quad (\text{by the definition of } \sim)\end{aligned}$$

- (b) $[(m, n)] + [(0, 0)] = [(m + 0, n + 0)] = [(m, n)]$. For the other relation, either argue that it holds by the first one and commutativity, or show directly that $[(0, 0)] + [(m, n)] = [(0 + m, 0 + n)] = [(m, n)]$.
- (c) $[(m, n)] + [(n, m)] = [(m + n, n + m)] = [(m + n, m + n)] = [(0, 0)]$ where the last relation holds since $(m + n, m + n) \sim (0, 0)$ as $m + n + 0 = m + n + 0$. The other relation holds since the first holds and addition is commutative.
- (d) Let us consider the case $m \geq n$ first. In this case, $m - n$ is a natural number and the relation $[(n, m)] = [(0, m - n)]$ is equivalent with $(n, m) \sim (0, m - n)$ and this last relation is, by definition of \sim equivalent with $n + m - n = m + 0$. This last relation is true since both $m + 0$ and $n + m - n$ are equal to m .

Let us consider the case $m < n$ now. In this case $n - m$ is a natural number and the relation $[(n, m)] = [(n - m, 0)]$ is equivalent with $(n, m) \sim (n - m, 0)$. This last relation is equivalent with $n + 0 = m + n - m$ by the definition of \sim . The relation $n + 0 = m + n - m$ is true since both sides are equal to n .

2. $[(6, 3)]$ can be shortened to $6 - 3 = 3$, $[(1000, 1005)]$ to $1000 - 1005 = -5$, $[(6, 8)]$ to $6 - 8 = -2$, $[(57, 56)]$ to $57 - 56 = 1$, and $[(56, 58)]$ to $56 - 58 = -2$. As $-5 < -2 = -2 < 1 < 3$, we have that

$$[(1000, 1005)] < [(6, 8)] = [(56, 58)] < [(57, 56)] < [(6, 3)].$$

3. $[(m, n)] \cdot [(n, m)] = [(mn, nm)] = [(mn, mn)] = [(1, 1)]$ where the last relation holds since $(mn, mn) \sim (1, 1)$ as $mn \cdot 1 = mn \cdot 1$. Similarly, $[(n, m)] \cdot [(m, n)] = [(nm, mn)] = [(nm, nm)] = [(1, 1)]$.
4. $[(5, 15)]$ can be shortened to $\frac{5}{15} = \frac{1}{3}$, $[(50, -100)]$ to $\frac{50}{-100} = \frac{-1}{2}$, $[(15, 10)]$ to $\frac{15}{10} = \frac{3}{2}$, $[(20, -10)]$ to $\frac{20}{-10} = -2$, and $[(10, 20)]$ to $\frac{10}{20} = \frac{1}{2}$. As $-2 < \frac{-1}{2} = \frac{-1}{2} < \frac{1}{3} < \frac{3}{2}$, we have that

$$[(20, -10)] < [(50, -100)] = [(-10, 20)] < [(5, 15)] < [(15, 10)].$$

5. Let $a = [(m, n)]$ and $b = [(k, l)]$ and assume that $ab = 0$ so that $[(mk, nl)] = [(0, 1)]$. This implies that $mk \cdot 1 = nl \cdot 0$ so that $mk = 0$. As \mathbb{Z} has no zero divisors, this implies that $m = 0$ or $k = 0$. If $m = 0$, then $a = [(0, n)] = [(0, 1)] = 0$. If $k = 0$, then $b = [(0, l)] = [(0, 1)] = 0$.

6. (a) Let a stand for the limit of this sequence in case it exists. Note that then $a = \lim_{n \rightarrow \infty} a_n$ and $a = \lim_{n \rightarrow \infty} a_{n+1}$ as well. To find the value of a let $n \rightarrow \infty$ in the equation $a_{n+1} = \sqrt{2 + a_n}$. The left side converges to a and the right side to $\sqrt{2 + a}$. So, a can be found from the equation $a = \sqrt{2 + a} \Rightarrow a^2 = 2 + a \Rightarrow a^2 - a - 2 = 0 \Rightarrow (a - 2)(a + 1) = 0 \Rightarrow a = 2$ or $a = -1$. Since -1 is an extraneous root (it does not satisfy the equation $a = \sqrt{2 + a}$), the limit of the sequence is $a = 2$. Alternatively, you can also argue that starting with the nonnegative term $a_0 = 0$, all the terms of the sequence are nonnegative and so the solution $a = -1$ can be discarded.
- (b) Let a stand for the limit of this sequence in case it exists. Note that then $a = \lim_{n \rightarrow \infty} a_n$ and $a = \lim_{n \rightarrow \infty} a_{n+1}$ as well. To find the value of a let $n \rightarrow \infty$ in the equation $a_{n+1} = \frac{1}{1 + a_n}$. The left side converges to a and the right side to $\frac{1}{1 + a}$. So, a can be found from the equation $a = \frac{1}{1 + a} \Rightarrow a(1 + a) = 1 \Rightarrow a^2 + a - 1 = 0 \Rightarrow a = \frac{-1 + \sqrt{5}}{2} \approx 0.618$ or $a = \frac{-1 - \sqrt{5}}{2} \approx -1.618$. Starting with the positive term $a_0 = 1$, all the terms of the sequence are positive, so the sequence converges towards the positive value $a = \frac{-1 + \sqrt{5}}{2} \approx 0.618$.


7. (a) $|(3, 5) \cup [8, 9]| = |(3, 5)| + |[8, 9]| = |\mathbb{R}| + |\mathbb{R}| = |\mathbb{R}|$ and $|(7, \infty)| = |\mathbb{R}|$.
- (b) Note that $[0, 9] \cup [7, \infty) = [0, \infty)$ and $(3, 5] \cup [0, \infty) = [0, \infty)$. So, $|(3, 5] \cup [0, 9] \cup [7, \infty)| = |[0, \infty)| = |\mathbb{R}|$ and $|(-\infty, 1]| = |\mathbb{R}|$.
- (c) Note that $\bigcup_{n \in \mathbb{N}} (-n, n) = (-1, 1) \cup (-2, 2) \cup (-3, 3) \cup \dots = (-\infty, \infty) = \mathbb{R}$. As $|(0, 1)| = |\mathbb{R}|$, the two sets have the same cardinality.
- (d) $\bigcap_{n \in \mathbb{N}} [0, n+1) = [0, 1) \cap [0, 2) \cap [0, 3) \cap \dots = [0, 1)$. As $|[0, 1)| = |\mathbb{R}|$, the two sets have the same cardinality.
- (e) $\bigcup_{n \in \mathbb{N}} (-\infty, -n) = (-\infty, 0) \cup (-\infty, -1) \cup (-\infty, -2) \cup (-\infty, -3) \cup \dots = (-\infty, 0)$. Since $|(-\infty, 0)| = |\mathbb{R}|$ and $|(1, \infty)| = |\mathbb{R}|$, the two sets have the same cardinality.
8. (a) $0.222222\dots = 0.2 + 0.02 + 0.002 + \dots = \frac{2}{10} + \frac{2}{10^2} + \frac{2}{10^3} + \dots = \sum_{n=1}^{\infty} 2 \left(\frac{1}{10}\right)^n$. Using the formula $\frac{ar^k}{1-r}$ with $a = 2$, $r = \frac{1}{10}$ and $k = 1$, we have that the sum is $\frac{\frac{2}{10}}{\frac{9}{10}} = \frac{2}{9}$.
- (b) $0.27272727\dots = 0.27 + 0.0027 + 0.000027 + \dots = \frac{27}{100} + \frac{27}{100^2} + \frac{27}{100^3} + \dots = \sum_{n=1}^{\infty} 27 \left(\frac{1}{100}\right)^n$. Using the formula $\frac{ar^k}{1-r}$ with $a = 27$, $r = \frac{1}{100}$ and $k = 1$, we have that the sum is $\frac{\frac{27}{100}}{\frac{99}{100}} = \frac{27}{99} = \frac{3}{11}$.
- (c) $1.2345454545\dots = 1.23 + 0.0045 + 0.000045 + 0.00000045 + \dots = 1.23 + \frac{45}{100^2} + \frac{45}{100^3} + \frac{45}{100^4} + \dots = 1.23 + \sum_{n=2}^{\infty} 45 \left(\frac{1}{100}\right)^n$. Using the formula $\frac{ar^k}{1-r}$ with $a = 45$, $r = \frac{1}{100}$ and $k = 2$, we have that the sum is $1.23 + \frac{\frac{45}{100^2}}{\frac{99}{100}} = \frac{123}{100} + \frac{45}{99(100)} = \frac{123(99)+45}{99(100)} = \frac{12222}{9900} = \frac{679}{550}$.
9. The complex number $-3i$ is on the negative part of y axis. Hence, $\theta = \frac{-\pi}{2}$. We have that $r = \sqrt{(-3)^2} = 3$.
- The complex number $\sqrt{2} - \sqrt{2}i$ is on the $y = -x$ line and in the fourth quadrant. Hence, $\theta = \frac{-\pi}{4}$. We have that $r = \sqrt{\sqrt{2}^2 + (-\sqrt{2})^2} = \sqrt{2+2}\sqrt{4} = 2$.
- The complex number $-\sqrt{3} + i$ is in the second quadrant. Hence, $\theta = \pi + \tan^{-1} \frac{1}{-\sqrt{3}} = \pi + \frac{-\pi}{6} = \frac{5\pi}{6}$. The modulus is $r = \sqrt{(-\sqrt{3})^2 + 1^2} = \sqrt{4} = 2$.
- The complex number $-2 - i$ is in the third quadrant. Hence, $\theta = \pi + \tan^{-1} \frac{-1}{-2} = \pi + \tan^{-1} \frac{1}{2} \approx \pi + 0.4636 \approx 3.605$. The modulus is $r = \sqrt{(-2)^2 + (-1)^2} = \sqrt{5} \approx 2.24$.
10. If $\theta = \frac{-\pi}{2}$, the number is on the negative part of y -axis. As $r = 5$, $(x, y) = (0, -5)$. Alternatively, $x = 5 \cos \frac{-\pi}{2} = 0$ and $y = 5 \sin \frac{-\pi}{2} = -5$.
- If $\theta = \frac{5\pi}{6}$ and $r = 2$, $x = r \cos \theta = 2 \cos \frac{5\pi}{6} = 2 \cdot \frac{-\sqrt{3}}{2} = -\sqrt{3}$ and $y = r \sin \theta = 2 \sin \frac{5\pi}{6} = 2 \cdot \frac{1}{2} = 1$. Thus, $(x, y) = (-\sqrt{3}, 1)$.
- If $\theta = \frac{-2\pi}{3}$ and $r = 3$, $x = r \cos \theta = 3 \cos \frac{-2\pi}{3} = 3 \cdot \frac{-1}{2} = \frac{-3}{2}$ and $y = r \sin \theta = 3 \sin \frac{-2\pi}{3} = 3 \cdot \frac{-\sqrt{3}}{2} = \frac{-3\sqrt{3}}{2}$. Thus, $(x, y) = (\frac{-3}{2}, \frac{-3\sqrt{3}}{2})$.
11. (a) From problem (1), we have that $z = -\sqrt{3} + i = 2e^{5\pi/6i}$. Hence, $z^4 = 2^4 e^{4 \cdot 5\pi/6i} = 16e^{10\pi/3i} = 16(\cos \frac{10\pi}{3} + i \sin \frac{10\pi}{3}) = 16(\frac{-1}{2} - \frac{\sqrt{3}}{2}i) = -8 - 8\sqrt{3}i$.
- (b) From problem (1), we have that $z = -2 - i \approx \sqrt{5}e^{3.605i}$. Hence, $z^6 \approx (\sqrt{5})^6 e^{6 \cdot 3.605i} = 125e^{21.63i} = 125(\cos 21.63 + i \sin 21.63) = 125(-0.936 + 0.352i) = -117 + 44i$.

12. (a) We need to find all five solutions of $z^5 = 32$. Note that 32 corresponds to the complex number $(32, 0)$ which is on the positive side of the x -axis so $\theta = 0$. The distance from $(32, 0)$ to the origin is 32 so $r = 32$. Hence, the five solutions of the characteristic equation can be found by the formula

$$\sqrt[5]{32}e^{\frac{0+2k\pi}{5}i} = 2e^{\frac{2k\pi}{5}i} \quad \text{for } k = 0, 1, \dots, 4.$$


These five solutions form a regular polygon with five sides on the circle of radius 2 centered at the origin.

$$\begin{aligned} k = 0 &\Rightarrow z_0 = 2e^{0i} = 2, \\ k = 1 &\Rightarrow z_1 = 2e^{\frac{2\pi}{5}i} = 2\left(\cos \frac{2\pi}{5} + i \sin \frac{2\pi}{5}\right) \approx 0.62 + 1.90i, \\ k = 2 &\Rightarrow z_2 = 2e^{\frac{4\pi}{5}i} = 2\left(\cos \frac{4\pi}{5} + i \sin \frac{4\pi}{5}\right) \approx -1.62 + 1.18i, \\ k = 3 &\Rightarrow z_3 = 2e^{\frac{6\pi}{5}i} = 2\left(\cos \frac{6\pi}{5} + i \sin \frac{6\pi}{5}\right) \approx -1.62 - 1.18i, \\ k = 4 &\Rightarrow z_4 = 2e^{\frac{8\pi}{5}i} = 2\left(\cos \frac{8\pi}{5} + i \sin \frac{8\pi}{5}\right) \approx 0.62 - 1.90i. \end{aligned}$$



- (b) $z^5 = -32 = 32e^{\pi i}$. Hence, $z_k = \sqrt[5]{32}e^{\frac{\pi+2k\pi}{5}i} = 2e^{\frac{(2k+1)\pi}{5}i}$ for $k = 0, 1, \dots, 4$.

$$\begin{aligned} k = 0 &\Rightarrow z_0 = 2e^{\frac{\pi}{5}i} = 2\left(\cos \frac{\pi}{5} + i \sin \frac{\pi}{5}\right) \approx 1.62 + 1.18i, \\ k = 1 &\Rightarrow z_1 = 2e^{\frac{3\pi}{5}i} = 2\left(\cos \frac{3\pi}{5} + i \sin \frac{3\pi}{5}\right) \approx -0.62 + 1.90i, \\ k = 2 &\Rightarrow z_2 = 2e^{\frac{5\pi}{5}i} = 2e^{\pi i} = 2(\cos \pi + i \sin \pi) = -2, \\ k = 3 &\Rightarrow z_3 = 2e^{\frac{7\pi}{5}i} = 2\left(\cos \frac{7\pi}{5} + i \sin \frac{7\pi}{5}\right) \approx -0.62 - 1.90i, \\ k = 4 &\Rightarrow z_4 = 2e^{\frac{9\pi}{5}i} = 2\left(\cos \frac{9\pi}{5} + i \sin \frac{9\pi}{5}\right) \approx 1.62 - 1.18i. \end{aligned}$$



- (c) $r = \sqrt{3^2 + 3^2} = \sqrt{18}$ or $3\sqrt{2}$, $\theta = \tan^{-1}(\frac{3}{3}) = \tan^{-1}(1) = \frac{\pi}{4}$, so $z = \sqrt[4]{18}e^{\pi/4i}$.


The four roots are obtained as

$$z_k = \sqrt[4]{\sqrt{18}}e^{\frac{\pi/4+2k\pi}{4}i} = 18^{1/8}e^{\frac{\pi+8k\pi}{16}i} \approx 1.435e^{\frac{\pi+8k\pi}{16}i}$$

for $k = 0, 1, 2, 3$. Thus,

$$\begin{aligned} z_0 &\approx 1.435e^{\pi/16i} = 1.435\left(\cos \frac{\pi}{16} + i \sin \frac{\pi}{16}\right) \approx 1.435(0.98 + i0.195) = 1.41 + 0.28i \\ z_1 &\approx 1.435e^{9\pi/16i} = 1.435\left(\cos \frac{9\pi}{16} + i \sin \frac{9\pi}{16}\right) \approx 1.435(-0.195 + 0.98i) = -0.28 + 1.41i \end{aligned}$$

$$\begin{aligned} z_2 &\approx 1.435e^{17\pi/16i} = 1.435\left(\cos \frac{17\pi}{16} + i \sin \frac{17\pi}{16}\right) = 1.435(-0.98 - 0.195i) = -1.41 - 0.28i \\ z_3 &\approx 1.435e^{25\pi/16i} = 1.435\left(\cos \frac{25\pi}{16} + i \sin \frac{25\pi}{16}\right) = 1.435(0.195 - 0.98i) = 0.28 - 1.41i \end{aligned}$$



- (d) $r = \sqrt{3^2 + (-3)^2} = \sqrt{18}$ or $3\sqrt{2}$, $\theta = \tan^{-1}(\frac{-3}{3}) = \tan^{-1}(-1) = \frac{-\pi}{4}$, so $z = \sqrt[4]{18}e^{-\pi/4i}$.

The four roots are obtained as

$$z_k = \sqrt[4]{\sqrt{18}} e^{-\pi/4 + 2k\pi i} = 18^{1/8} e^{-\pi + 8k\pi i} \approx 1.435 e^{-\pi + 8k\pi i}$$


for $k = 0, 1, 2, 3$. Thus,

$$z_0 \approx 1.435 e^{-\pi/16i} = 1.435 \left(\cos \frac{\pi}{16} + i \sin \frac{\pi}{16} \right) \approx 1.435 (0.98 - i0.195) = 1.41 - 0.28i$$

$$z_1 \approx 1.435 e^{7\pi/16i} = 1.435 \left(\cos \frac{7\pi}{16} + i \sin \frac{7\pi}{16} \right) \approx 1.435 (0.195 + i0.98) = 0.28 + 1.41i$$

$$z_2 \approx 1.435 e^{15\pi/16i} = 1.435 \left(\cos \frac{15\pi}{16} + i \sin \frac{15\pi}{16} \right) = 1.435 (-0.98 + 0.195i) = -1.41 + 0.28i$$

$$z_3 \approx 1.435 e^{23\pi/16i} = 1.435 \left(\cos \frac{23\pi}{16} + i \sin \frac{23\pi}{16} \right) = 1.435 (-0.195 - 0.98i) = -0.28 - 1.41i$$



13. (a) $\cos z + i \sin z = \frac{1}{2}(e^{iz} + e^{-iz}) + i \frac{1}{2i}(e^{iz} - e^{-iz}) = \frac{1}{2}(e^{iz} + e^{-iz} + e^{iz} - e^{-iz}) = \frac{1}{2}(2e^{iz}) = e^{iz}$.
- (b) $\sin^2 z + \cos^2 z = \frac{-1}{4}(e^{iz} - e^{-iz})^2 + \frac{1}{4}(e^{iz} + e^{-iz})^2 = \frac{-1}{4}(e^{2iz} - 2 + e^{-2iz}) + \frac{1}{4}(e^{2iz} + 2 + e^{-2iz}) = \frac{1}{4}(-e^{2iz} + 2 - e^{-2iz} + e^{2iz} + 2 + e^{-2iz}) = \frac{1}{4}(4) = 1$.
- (c) $\sin^2 z = \left(\frac{1}{2i}(e^{iz} - e^{-iz}) \right)^2 = \frac{1}{-4}(e^{2iz} - 2 + e^{-2iz}) = \frac{1}{4}(2 - (e^{2iz} + e^{-2iz})) = \frac{1}{2} \left(1 - \frac{1}{2}(e^{2iz} + e^{-2iz}) \right) = \frac{1}{2}(1 - \cos(2z))$
- (d) $\cos^2 z = \left(\frac{1}{2}(e^{iz} + e^{-iz}) \right)^2 = \frac{1}{4}(e^{2iz} + 2 + e^{-2iz}) = \frac{1}{4}(2 + e^{2iz} + e^{-2iz}) = \frac{1}{2} \left(1 + \frac{1}{2}(e^{2iz} + e^{-2iz}) \right) = \frac{1}{2}(1 + \cos(2z))$